



## RESEARCH ARTICLE

### ON ANEW LAGRANGIAN MECHANICAL SYSTEMS ON GENERALIZED-QUATERNIONIC Kähler MANIFOLDS

Gebreel Mohammed Khur Baba Gebreel , Eltayeb Awad Adam and  
Ibrahim Yousif Ibrahim Abad Alrhman

Department of Mathematics, Faculty of Education, West Kordufan University, Khartoum-Sudan

#### ARTICLE INFO

##### Article History:

Received 25<sup>th</sup> November, 2025  
Received in revised form  
20<sup>th</sup> December, 2025  
Accepted 18<sup>th</sup> January, 2026  
Published online 27<sup>th</sup> February, 2026

##### Keywords:

Preliminaries; Generalized-Quaternionic  
Kähler geometry; Lagrangian Mechanical  
Systems.

\*Corresponding author: Bheem Rao, T.

Copyright©2026, Bheem Rao and Anil Kumar. 2026. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Bheem Rao, T. and Anil Kumar, V. 2026. "On Anew Lagrangian Mechanical Systems On Generalized-Quaternionic Kähler Manifolds". *International Journal of Current Research*, 18, (02), 36190-36194.

#### ABSTRACT

In this paper, we study Lagrangian mechanical system on generalized-quaternionic Kähler Manifolds by using vector field:  $Y = \sum_{a=0}^3 Y^{an+} \frac{\partial}{\partial x_{an+i}}$  in a canonical local basis  $\{G\}$  of  $V$ . Finally, the geometrical – Physical results related to generalized-quaternionic Kähler mechanical systems are also given .

## INTRODUCTION

Modern differential geometry plays an important role to explain the dynamics of Lagrangians. So , if  $Q$  is an  $m - dimensional$  configuration manifold and  $L : TQ \rightarrow R$  is a regular Lagrangian function , then it is well-known that there is a unique vector field  $\xi$  on  $TQ$  such that the dynamic equation is given by

$$i_{\xi}\Phi_L = dE_L \rightarrow \tag{1}$$

Where  $\Phi_L = -dd_JL$  indicates the symplectic form ,  $J : TQ \rightarrow TQ$  respectively an almost tangent or complex or paracomplex structure such that  $JoJ = 0$  or  $JoJ = -1$  or  $JoJ = 1$ . Also  $E_L = V(L) - L$  is an energy function and  $V = J(\xi)$  a Liouville vector field. The triple  $(TQ, \Phi_L, \xi)$  is called Lagrangian system on the tangent bundle. Also, modern differential geometry provides a good framework in which develops the dynamics of Hamiltonians. Therefore, if  $Q$  is an  $m$ -dimensional configuration manifold and  $H : T^*Q \rightarrow R$  is a regular Hamiltonian function, then there is a unique vector field  $X$  on  $T^*Q$  such that the dynamic equation is given by :

$$i_X\Phi = dH \rightarrow \tag{2}$$

Where  $\Phi$  indicates the canonical symplectic form. The triple  $(T^*Q, \Phi, X)$  is called Hamiltonian system on the cotangent bundle  $T^*Q$ . Nowadays , there are many studies about Lagrangian and Hamiltonian dynamics, mechanics, formalisms, Systems and equations ( 1, 2 ) and therein. There are real, complex, paracomplex and other analogues. As known it is possible to produce different analogues in different spaces. Quaternions were invented by Sir William Rowan Hamiltonian as an extension to the complex numbers. Hamiltonian's defining relation is most succinctly written as:

$$i^2 = j^2 = k^2 = -1, ijk = -1.$$

Split quaternions are given by

$$i^2 = -1, j^2 = 1 = k^2, ijk = 1$$

Generalized-quaternions are defined as

$$i^2 = -a, j^2 = -b, k^2 = -ab, ijk = -ab$$

If it is compared to the calculus of vectors, quaternions have slipped into the realm of obscurity. They do however still find use in the computation of rotations. A lot of physical laws in classical, relativistic, and quantum mechanics can be written pleasantly by means of quaternions. Some physicists hope they will find deeper understanding of the universe by restating basis principles in terms of quaternion algebra ( 3, 4).

**Preliminaries:** In this paper, we present equations related to Lagrangian mechanical systems on a generalized-quaternionic kähler manifold, all mathematical objects and mappings are assumed to be smooth, i.e. infinitely differentiable and Einstein convention of summarizing is adopted.  $\mathcal{F}(M), \mathcal{K}(M)$  and  $\Lambda^1(M)$  denote the set of functions on M, the set of vector fields on M and the set of 1-forms on M, respectively.

**Theorem**

Let  $f$  be differentiable  $\phi, \psi$  are 1-forme, then ( 5 , 6 ):

- $d(f\phi) = df \wedge \phi + f d\phi$
- $d(\phi \wedge \psi) = d\phi \wedge \psi - \phi \wedge d\psi$

**Definition ( Kronecker’s delta )**

A kronecker’s delta denote by  $\delta$  and defined as follows ( 5 ):

$$\delta_i^j = \begin{cases} 1; & \text{if } i = j \\ 0; & \text{if } i \neq j \end{cases}$$

**Generalized-Quaternionic kähler Manifolds:** A generalized almost quaternion structure on the manifold M is a sub bundle of the bundle of endomorphisms of the tangent bundle of M, whose standard fiber is the algebra of generalized-quaternions. A generalized almost quaternion structure on a pseudo-Riemannian manifold is called a generalized-quaternion Hermitian if the following conditions hold:

( i ) The endomorphisms  $F, G$  and  $H$  of  $T_x M$  satisfy

$$F^2 = -aI, G^2 = -bI, H^2 = -abI, FG = H, GH = bF, HF = aG \rightarrow \tag{3}$$

( ii ) The compatibility equations are given by, for  $X, Y \in T_x M$

$$g(FX, FY) = ag(X, Y), g(GX, GY) = bg(X, Y), g(HX, HY) = abg(X, Y) \rightarrow \tag{4}$$

Where I denotes the identity tensor of type(1,1) in M. In particular, 2-form  $Q$  defined by  $Q(X, Y) = (X, FY) = (X, GY) = (X, HY)$  on M is called the Kähler form of the endomorphisms  $F, G$  and  $H$ . If the Kähler form  $Q$  on M is closed, i.e.  $dQ = 0$ , the manifold M is called a generalized-quaternionic Kähler manifold (for detail see ( 7 ) ). If  $a = b = 1, M$  is quaternion manifold. If  $a = 1, b = -1, M$  is para-quaternion manifold. The bundle V is asset that locally admits a basis  $\{F, G, H\}$  satisfying ( 1 ) and ( 2 ) in any coordinate neighborhood  $U \subset M$  such that  $M = \cup U$  (8). Then  $V$  is called a generalized-quaternionic structure in M. The pair  $(M, V)$  denotes a generalized-quaternionic manifold with  $V$ . The structure  $V$  with such a Riemannian metric  $g$  is called a generalized-quaternionic metric structure. The triple  $(M, g, V)$  denotes a generalized-quaternion metric manifold.

Let  $\{x_i, x_{n+i}, x_{2n+i}, x_{3n+i}\}, i = \overline{1, n}$  be a real coordinate system on a neighborhood  $U$  of  $M$ , and let  $\left\{ \frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_{n+i}}, \frac{\partial}{\partial x_{2n+i}}, \frac{\partial}{\partial x_{3n+i}} \right\}$  and  $\{dx_i, dx_{n+i}, dx_{2n+i}, dx_{3n+i}\}$  be natural bases over  $R$  of the tangent space  $T(M)$  and the cotangent space  $T^*(M)$  of  $M$ , respectively. Taking into consideration ( 1 ), then we can obtain the expression as follows:

$$\begin{aligned} F\left(\frac{\partial}{\partial x_i}\right) &= a \frac{\partial}{\partial x_{n+i}}, G\left(\frac{\partial}{\partial x_i}\right) = -b \frac{\partial}{\partial x_{2n+i}}, H\left(\frac{\partial}{\partial x_i}\right) = -ab \frac{\partial}{\partial x_{3n+i}} \\ F\left(\frac{\partial}{\partial x_{n+i}}\right) &= -a \frac{\partial}{\partial x_i}, G\left(\frac{\partial}{\partial x_{n+i}}\right) = b \frac{\partial}{\partial x_{3n+i}}, H\left(\frac{\partial}{\partial x_{n+i}}\right) = -ab \frac{\partial}{\partial x_{2n+i}} \rightarrow \\ F\left(\frac{\partial}{\partial x_{2n+i}}\right) &= a \frac{\partial}{\partial x_{3n+i}}, G\left(\frac{\partial}{\partial x_{2n+i}}\right) = -b \frac{\partial}{\partial x_i}, H\left(\frac{\partial}{\partial x_{2n+i}}\right) = -ab \frac{\partial}{\partial x_{n+i}} \\ F\left(\frac{\partial}{\partial x_{3n+i}}\right) &= -a \frac{\partial}{\partial x_{2n+i}}, G\left(\frac{\partial}{\partial x_{3n+i}}\right) = b \frac{\partial}{\partial x_{n+i}}, H\left(\frac{\partial}{\partial x_{3n+i}}\right) = -ab \frac{\partial}{\partial x_i} \end{aligned} \tag{5}$$

A canonical local basis  $\{F^*, G^*, H^*\}$  of  $V^*$  of the cotangent space  $T^*(M)$  of a manifold  $M$  satisfies the condition as follows:

$$F^{*2} = -aI, G^{*2} = -bI, H^{*2} = -abI, F^*G^* = H^*, G^*H^* = bF, H^*F^* = aG^* \rightarrow$$

Defining by

$$F^*(dx_i) = adx_{n+i}, G^*(dx_i) = -bdx_{2n+i}, H^*(dx_i) = -abdx_{3n+i}$$

$$F^*(dx_{n+i}) = -adx_i, G^*(dx_{n+i}) = bdx_{3n+i}, H^*(dx_{n+i}) = -abdx_{2n+i} \rightarrow$$

$$F^*(dx_{2n+i}) = adx_{3n+i}, G^*(dx_{2n+i}) = -bdx_i, H^*(dx_{2n+i}) = -abdx_{n+i}$$

$$F^*(dx_{3n+i}) = -adx_{2n+i}, G^*(dx_{3n+i}) = bdx_{n+i}, H^*(dx_{3n+i}) = -abdx_i$$

**Lagrangian Mechanical Systems :** Here, we obtain Euler-Lagrange equation for quantum and classical mechanics by means of a canonical local basis  $\{F, G, H\}$  of  $V$  on a generalized-quaternionic *kähler* manifold  $(M, g, V)$ .

First:

$$a \frac{\partial}{\partial t} \left( \frac{\partial L}{\partial x_i} \right) + \frac{\partial L}{\partial x_{n+i}} = 0, a \frac{\partial}{\partial t} \left( \frac{\partial L}{\partial x_{n+i}} \right) - \frac{\partial L}{\partial x_i} = 0,$$

$$a \frac{\partial}{\partial t} \left( \frac{\partial L}{\partial x_{2n+i}} \right) + \frac{\partial L}{\partial x_{3n+i}} = 0, a \frac{\partial}{\partial t} \left( \frac{\partial L}{\partial x_{3n+i}} \right) - \frac{\partial L}{\partial x_{2n+i}} = 0 \rightarrow$$

Second, let  $G$  take a local basis element on the generalized-quaternionic *kähler* manifold  $(M, g, V)$ , and  $\{x_i, x_{n+i}, x_{2n+i}, x_{3n+i}\}$  be its coordinate functions. Let semispray be the vector field  $Y$  determined by

$$Y = Y^i \frac{\partial}{\partial x_i} + Y^{n+i} \frac{\partial}{\partial x_{n+i}} + Y^{2n+i} \frac{\partial}{\partial x_{2n+i}} + Y^{3n+i} \frac{\partial}{\partial x_{3n+i}} \rightarrow$$

Where  $Y^i = \dot{x}_i, Y^{n+i} = \dot{x}_{n+i}, Y^{2n+i} = \dot{x}_{2n+i}, Y^{3n+i} = \dot{x}_{3n+i}$ .

This equation ( 9 ) can be written concise manner:

$$Y = \sum_{a=0}^3 Y^{an+i} \frac{\partial}{\partial x_{an+i}} \rightarrow$$

And the dot indicates the derivative with respect to time  $t$ . The vector field defined by:

$$V_G = G(Y) = -bY^i \frac{\partial}{\partial x_{2n+i}} + bY^{n+i} \frac{\partial}{\partial x_{3n+i}} - bY^{2n+i} \frac{\partial}{\partial x_i} + bY^{3n+i} \frac{\partial}{\partial x_{n+i}} : \mathcal{F}(M) \rightarrow \Lambda^1 M \rightarrow$$

Is named a Liouville vector field on the generalized-quaternionic *kähler* manifold  $(M, g, V)$ , for  $G$ , the closed generalized-quaternionic *kähler* form is the closed 2-form given by  $\Phi_L^G = -dd_G L$  such that

$$d = \frac{\partial}{\partial x_i} dx_i + \frac{\partial}{\partial x_{n+i}} dx_{n+i} + \frac{\partial}{\partial x_{2n+i}} dx_{2n+i} + \frac{\partial}{\partial x_{3n+i}} dx_{3n+i} \rightarrow$$

This equation (12) can be written concise manner

$$d = \sum_{a=0}^3 \frac{\partial}{\partial x_{an+i}} dx_{an+i} \rightarrow$$

$$d_G L = -b \frac{\partial L}{\partial x_{2n+i}} dx_i + b \frac{\partial L}{\partial x_{3n+i}} dx_{n+i} - b \frac{\partial L}{\partial x_i} dx_{2n+i} + b \frac{\partial L}{\partial x_{n+i}} dx_{3n+i} : \mathcal{F}(M) \rightarrow \Lambda^1 M \rightarrow$$

Then we have:

$$\Phi_L^G = b \frac{\partial^2 L}{\partial x_j \partial x_{2n+i}} dx_j \wedge dx_i - b \frac{\partial^2 L}{\partial x_j \partial x_{3n+i}} dx_j \wedge dx_{n+i} + b \frac{\partial^2 L}{\partial x_j \partial x_i} dx_j \wedge dx_{2n+i}$$

$$- b \frac{\partial^2 L}{\partial x_j \partial x_{n+i}} dx_j \wedge dx_{3n+i} + b \frac{\partial^2 L}{\partial x_{n+j} \partial x_{2n+i}} dx_{n+j} \wedge dx_i - b \frac{\partial^2 L}{\partial x_{n+j} \partial x_{3n+i}} dx_{n+j} \wedge dx_{n+i}$$

$$+ b \frac{\partial^2 L}{\partial x_{n+j} \partial x_i} dx_{n+j} \wedge dx_{2n+i} - b \frac{\partial^2 L}{\partial x_{n+j} \partial x_{n+i}} dx_{n+j} \wedge dx_{3n+i} + b \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{2n+i}} dx_{2n+j} \wedge dx_i$$

$$- b \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{3n+i}} dx_{2n+j} \wedge dx_{n+i} + b \frac{\partial^2 L}{\partial x_{2n+j} \partial x_i} dx_{2n+j} \wedge dx_{2n+i} - b \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{n+i}} dx_{2n+j} \wedge dx_{3n+i}$$

$$+ b \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{2n+i}} dx_{3n+j} \wedge dx_i - b \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{3n+i}} dx_{3n+j} \wedge dx_{n+i} + b \frac{\partial^2 L}{\partial x_{3n+j} \partial x_i} dx_{3n+j} \wedge dx_{2n+i}$$

$$- b \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{n+i}} dx_{3n+j} \wedge dx_{3n+i}$$

$$\therefore i_Y = Y^i \frac{\partial}{\partial x_i} + Y^{n+i} \frac{\partial}{\partial x_{n+i}} + Y^{2n+i} \frac{\partial}{\partial x_{2n+i}} + Y^{3n+i} \frac{\partial}{\partial x_{3n+i}}$$

Then we calculate

$$\begin{aligned} i_Y \Phi_L^G &= bY^i \frac{\partial^2 L}{\partial x_j \partial x_{2n+i}} \delta_i^j dx_i - bY^i \frac{\partial^2 L}{\partial x_j \partial x_{2n+i}} dx_j - bY^i \frac{\partial^2 L}{\partial x_j \partial x_{3n+i}} \delta_i^j dx_{n+i} + bY^{n+i} \frac{\partial^2 L}{\partial x_j \partial x_{3n+i}} dx_j \\ &+ bY^i \frac{\partial^2 L}{\partial x_j \partial x_i} \delta_i^j dx_{2n+i} - bY^{2n+i} \frac{\partial^2 L}{\partial x_j \partial x_i} dx_j - bY^i \frac{\partial^2 L}{\partial x_j \partial x_{n+i}} \delta_i^j dx_{3n+i} + bY^{3n+i} \frac{\partial^2 L}{\partial x_j \partial x_{n+i}} dx_j \\ &+ bY^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{2n+i}} \delta_{n+i}^{n+j} dx_i - bY^i \frac{\partial^2 L}{\partial x_{n+j} \partial x_{2n+i}} dx_{n+j} - bY^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{3n+i}} \delta_{n+i}^{n+j} dx_{n+i} \\ &+ bY^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{3n+i}} dx_{n+j} + bY^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_i} \delta_{n+i}^{n+j} dx_{2n+i} - bY^{2n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_i} dx_{n+j} \\ &- bY^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{n+i}} \delta_{n+i}^{n+j} dx_{3n+i} + bY^{3n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{n+i}} dx_{n+j} + bY^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{2n+i}} \delta_{2n+i}^{2n+j} dx_i \\ &- bY^i \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{2n+i}} dx_{2n+j} - bY^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{3n+i}} \delta_{2n+i}^{2n+j} dx_{n+i} + bY^{n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{3n+i}} dx_{2n+j} \\ &+ bY^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_i} \delta_{2n+i}^{2n+j} dx_{2n+i} - bY^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_i} dx_{2n+j} - bY^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{n+i}} \delta_{2n+i}^{2n+j} dx_{3n+i} \\ &+ bY^{3n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{n+i}} dx_{2n+j} + bY^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{2n+i}} \delta_{3n+i}^{3n+j} dx_i - bY^i \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{2n+i}} dx_{3n+j} \\ &- bY^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{3n+i}} \delta_{3n+i}^{3n+j} dx_{n+i} + bY^{n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{3n+i}} dx_{3n+j} + bY^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_i} \delta_{3n+i}^{3n+j} dx_{2n+i} \\ &- bY^{2n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_i} dx_{3n+j} - bY^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{n+i}} \delta_{3n+i}^{3n+j} dx_{3n+i} + bY^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{n+i}} dx_{3n+j} \end{aligned}$$

The Energy function is

$$\begin{aligned} E_L^G &= V_G(L) - L \\ E_L^G &= -bY^i \frac{\partial L}{\partial x_{2n+i}} + bY^{n+i} \frac{\partial L}{\partial x_{3n+i}} - bY^{2n+i} \frac{\partial L}{\partial x_i} + bY^{3n+i} \frac{\partial L}{\partial x_{3n+i}} - L \rightarrow \end{aligned} \tag{15}$$

$$\begin{aligned} dE_L^G &= -bY^i \frac{\partial^2 L}{\partial x_j \partial x_{2n+i}} dx_j + bY^{n+i} \frac{\partial^2 L}{\partial x_j \partial x_{3n+i}} dx_j - bY^{2n+i} \frac{\partial^2 L}{\partial x_j \partial x_i} dx_j + bY^{3n+i} \frac{\partial^2 L}{\partial x_j \partial x_{n+i}} dx_j \\ &- bY^i \frac{\partial^2 L}{\partial x_{n+j} \partial x_{2n+i}} dx_{n+j} + bY^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{3n+i}} dx_{n+j} - bY^{2n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_i} dx_{n+j} + bY^{3n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{n+i}} dx_{n+j} \\ &- bY^i \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{2n+i}} dx_{2n+j} + bY^{n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{3n+i}} dx_{2n+j} - bY^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_i} dx_{2n+j} \\ &+ bY^{3n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{n+i}} dx_{2n+j} - bY^i \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{2n+i}} dx_{3n+j} + bY^{n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{3n+i}} dx_{3n+j} \\ &- bY^{2n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_i} dx_{3n+j} + bY^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{n+i}} dx_{3n+j} - \frac{\partial L}{\partial x_j} dx_j - \frac{\partial L}{\partial x_{n+j}} dx_{n+j} \\ &- \frac{\partial L}{\partial x_{2n+j}} dx_{2n+j} - \frac{\partial L}{\partial x_{3n+j}} dx_{3n+j} \end{aligned}$$

Using equation (1) and also considering an Integral curve of Y , we obtain the equation given by:

$$i_Y \Phi_L^G = dE_L^G$$

From which we get

$$\begin{aligned} &(bY^i \frac{\partial^2 L}{\partial x_j \partial x_{2n+i}} \delta_i^j dx_i + bY^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{2n+i}} \delta_{n+i}^{n+j} dx_i + bY^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{2n+i}} \delta_{2n+i}^{2n+j} dx_i + \\ &bY^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{2n+i}} \delta_{3n+i}^{3n+j} dx_i) - (bY^i \frac{\partial^2 L}{\partial x_j \partial x_{3n+i}} \delta_i^j dx_{n+i} + bY^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{3n+i}} \delta_{n+i}^{n+j} dx_{n+i} \\ &+ bY^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{3n+i}} \delta_{2n+i}^{2n+j} dx_{n+i} + bY^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{3n+i}} \delta_{3n+i}^{3n+j} dx_{n+i}) + (bY^i \frac{\partial^2 L}{\partial x_j \partial x_i} \delta_i^j dx_{2n+i} \\ &+ bY^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_i} \delta_{n+i}^{n+j} dx_{2n+i} + bY^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_i} \delta_{2n+i}^{2n+j} dx_{2n+i} + bY^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_i} \delta_{3n+i}^{3n+j} dx_{2n+i}) \\ &- (bY^i \frac{\partial^2 L}{\partial x_j \partial x_{n+i}} \delta_i^j dx_{3n+i} + bY^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{n+i}} \delta_{n+i}^{n+j} dx_{3n+i} + bY^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{n+i}} \delta_{2n+i}^{2n+j} dx_{3n+i} \\ &+ bY^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{n+i}} \delta_{3n+i}^{3n+j} dx_{3n+i}) = -(\frac{\partial L}{\partial x_j} dx_j + \frac{\partial L}{\partial x_{n+j}} dx_{n+j} + \frac{\partial L}{\partial x_{2n+j}} dx_{2n+j} + \frac{\partial L}{\partial x_{3n+j}} dx_{3n+j}) \end{aligned}$$

$$\text{If } i = j \Rightarrow \delta_i^j = 1, \delta_{n+i}^{n+i} = 1, \delta_{2n+i}^{2n+i} = 1, \delta_{3n+i}^{3n+i} = 1.$$

And therefore

$$\begin{aligned}
& b \left[ Y^i \frac{\partial^2 L}{\partial x_j \partial x_{2n+i}} + Y^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{2n+i}} + Y^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{2n+i}} + Y^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{2n+i}} \right] dx_i - b \left( Y^i \frac{\partial^2 L}{\partial x_j \partial x_{3n+i}} \right. \\
& + Y^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{3n+i}} + Y^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{3n+i}} + Y^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{3n+i}} \left. \right) dx_{n+j} + b \left( Y^i \frac{\partial^2 L}{\partial x_j \partial x_i} + Y^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_i} \right. \\
& + Y^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_i} + Y^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_i} \left. \right) dx_{2n+j} - b \left( Y^i \frac{\partial^2 L}{\partial x_j \partial x_{n+i}} + Y^{n+i} \frac{\partial^2 L}{\partial x_{n+j} \partial x_{n+i}} + Y^{2n+i} \frac{\partial^2 L}{\partial x_{2n+j} \partial x_{n+i}} \right. \\
& \left. + Y^{3n+i} \frac{\partial^2 L}{\partial x_{3n+j} \partial x_{n+i}} \right) dx_{3n+i} + \left[ \frac{\partial L}{\partial x_j} dx_j + \frac{\partial L}{\partial x_{n+j}} dx_{n+j} + \frac{\partial L}{\partial x_{2n+j}} dx_{2n+j} + \frac{\partial L}{\partial x_{3n+j}} dx_{3n+j} \right] = 0 \rightarrow (16)
\end{aligned}$$

In this equation can be concise manner

$$\begin{aligned}
& b \sum_{a=0}^3 Y^{an+i} \frac{\partial^2 L}{\partial x_{an+j} \partial x_{2n+i}} dx_i + \frac{\partial L}{\partial x_j} dx_j - b \sum_{a=0}^3 Y^{an+i} \frac{\partial^2 L}{\partial x_{an+j} \partial x_{3n+i}} dx_{n+i} + \frac{\partial L}{\partial x_{n+j}} dx_{n+j} \\
& + b \sum_{a=0}^3 Y^{an+i} \frac{\partial^2 L}{\partial x_{an+j} \partial x_i} dx_{2n+i} + \frac{\partial L}{\partial x_{2n+j}} dx_{2n+j} - b \sum_{a=0}^3 Y^{an+i} \frac{\partial^2 L}{\partial x_{an+j} \partial x_{n+i}} dx_{3n+i} \\
& + \frac{\partial L}{\partial x_{3n+j}} dx_{3n+j} = 0 \rightarrow (17)
\end{aligned}$$

Then we have the equations

$$\begin{aligned}
& b \frac{\partial}{\partial t} \left( \frac{\partial L}{\partial x_i} \right) + \frac{\partial L}{\partial x_{2n+i}} = 0, b \frac{\partial}{\partial t} \left( \frac{\partial L}{\partial x_{n+i}} \right) - \frac{\partial L}{\partial x_{3n+i}} = 0 \\
& b \frac{\partial}{\partial t} \left( \frac{\partial L}{\partial x_{2n+i}} \right) + \frac{\partial L}{\partial x_i} = 0, b \frac{\partial}{\partial t} \left( \frac{\partial L}{\partial x_{3n+i}} \right) - \frac{\partial L}{\partial x_{n+i}} = 0 \rightarrow (18)
\end{aligned}$$

Thus equations obtained in Eq (18) are called Euler-Lagrange equations constructed by means of  $\Phi_L^G$  on a generalized-quaternionic *kähler* manifold  $(M, g, V)$  and then triples  $(M, \Phi_L^G, Y)$  are named Mechanical systems on a generalized-quaternionic *kähler* manifold  $(M, g, V)$ .

## REFERENCES

- De Leon M. and P. R. Rodrigues, Methods of Differential Geometry in Analytical Mechanics, North-Holland Mathematics Studies, Vol. 152 ( Elsevier, Amsterdam, 1989 ).
- Tekkoyun, M. Mechanical systems on para-quaternionic *Kähler* manifolds, XVI. Geometrical Seminar, 20-25 September, Vrnjacka Banja, Serbia, 2009.
- Stahlke, D. Quaternions in classical mechanics, Phys. 621. <http://www.stahlke.org/dan/phys-papers/quaternion-paper.pdf>.
- Jafari M. and Y. Yayli, Generalized quaternions and their algebraic properties, submitted for publication.
- Gebreel Mohammed Khur Baba, M.A. Bashir . On a new dynamics On *Kähler* manifold using Canonical basis, International Journal of information research and review . vol. 04, Issue, 02 ,pp.3713-3722 ,February, 2017.
- Gebreel Mohammed Khur Baba, M.A. Bashir . On the Standard Cliffordian Hamiltonian Formulation of Symplectic Mechanics using Frame and co-Frame Fields, Global Journal of Science frontier research, Volume 20 Issue1. 2020.
- Arsen'eva, O. E. On the geometry of quaternion-Hermitian manifolds, Russian Math. Surveys 48 (5) (1993 ) 159.
- Dancer, A. S. H. R. Jorgensen and A. F. Swann, Metric geometries over the split quaternions, preprint (2004) , arXiv:0412.215.
- Tekkoyun, M. Yusuf Yayli , 2011, Mechanical systems on Generalized-quaternionic *kähler* manifold , International Journal of Geometric Methods in Modern Physics vol.8 , No.7(2011)1419-1431, © Word Scientific publishing Company , DOI:10.1142/S02198811005804.

\*\*\*\*\*