



RESEARCH ARTICLE

EPIDEMIOLOGICAL STUDIES OF HELMINTH INFECTED FISHES OF SHALLABUGH WETLAND

*Ibraq Khurshid and Fayaz Ahmad

Department of Zoology, University of Kashmir, Srinagar-190 006, Kashmir

ARTICLE INFO

Article History:

Received 26th March, 2013
Received in revised form
18th April, 2013
Accepted 27th May, 2013
Published online 15th June, 2013

Key words:

Fish; Parasite;
Helminth; *Schizothorax* spp,
Cyprinus spp. Wetland.

ABSTRACT

Helminths are of great interest to the ecologists because of their ordered life cycle. They are also considered as one of the important sensitive parasites to any changes in the water quality. Helminth parasites of exotic and indigenous fish were examined with the aim to evaluate their possible use as biological tags. This study quantified the seasonal prevalence and intensity of the helminth parasites in Shallabugh Wetland. Fish hosts studied includes two native fish of genus - *Schizothorax* and two exotic fish of genus - *Cyprinus*. 486 Fish specimens were examined out of which 197 were infected showing the overall prevalence of 40.53%. Our results show highest prevalence of helminth parasites in all the hosts examined viz., *S. niger* =59.25%, *S. labiatus* = 55.55%, *C. c. communis* = 61.11% and *C. c. specularis* = 55.55%, during summer months than spring and autumn. Positive relation between the infection indices of Helminth parasites and temperature suggests that helminth parasites can be used as stress indicators. It is further suggested that helminth parasites should be integrated with biomonitoring programmes, as they can provide supplementary information on pollution.

Copyright, IJCR, 2013, Academic Journals. All rights reserved.

INTRODUCTION

Helminths are a major cause of reduced productivity in fishes, characterised by devastating effects on fish health in terms of mortality and morbidity, particularly in developing countries. Various risk factors, including those of the host and environment, play an important role in the onset of helminth infection in fishes. The effect of the helminth infections on production of particular fish spp. depends mostly on the age of the fish, genotype, parasite species involved and the intensity of helminths. The climate in a particular locality is also one of the important factors that determine the type severity of parasitic infections in fishes. Fishing is an economically important aspect of the people in Kashmir valley; as the most people live in rural areas earn their live hood through fishing. Therefore, it is of prime importance to identify the helminths and to demonstrate the associated factors that influence the epidemiology of helminth infection of fishes in this climatically different and unique zone of the world.

MATERIALS AND METHODS

Fishes used for the study were bought from same fisherman operating with gill nets and cast nets. The fish samples were kept in plastic coolers containing river water and ice blocks before being transported to the laboratory. In the laboratory the fishes were identified to species level using keys provided by Holden and Reed (1972) and Lowe-Mc Connel (1972). Fish standard length (SL – from the snout to the base of the caudal peduncle) were determined with a meter rule while body weight (BW) was determined using a weighing balance. The gills, muscles, intestine, stomach and oesophagus of the fish were examined for parasites. Parasites recovered from each site were properly washed, fixed in alcohol-formol- acetic acid according to Olurin and Somorin (2006) and site of infection noted. Identification of parasites was carried out according to Yamaguti (1958, 1959, 1961 and 1963).

*Corresponding author: Ibraq Khurshid
Department of Zoology, University of Kashmir, Srinagar-190 006,
Kashmir

Data analysis

Percentages (%) to measure prevalence and chi-square to measure association between the parasitism and age, sex, breed and season were the statistical tools applied. The data was analyzed using Statistical packages SPSS software version 17. In all the analyses, confidence level was held at 95% and $P < 0.005$ was set for significance.

RESULTS

During the course of study, a total of 486 fish samples were subjected to parasitological investigation. Out of these 197 fish were found infected with 507 parasites. Thus the overall prevalence of parasitic infection in these fishes was 40.53%. Parasites recovered were Acanthocephalans, Cestodes and Trematodes. The gills, stomach and intestine were infected; no parasites were found infecting the liver and eye lens.

Seasonal prevalence

The data pooled for seasonal estimation of helminth infection revealed definite seasonality of infection ($p < 0.05$) in fishes over a period of 1 year, with highest infection in summer and lowest in winter. (Table 1). The maximum prevalence of helminth infection (59.25) occurred in summer season, and they were lowest in winter (31.81) with significant difference between the seasons ($p > 0.05$). The rainy season which started in spring and continued to early summers, made the environmental conditions more favourable for the development and survival of the pre-parasitic stages of helminth parasites and lead to an increased availability of infective stages in rainy and post-rainy season. This resulted in higher prevalence and level of parasitism in summer and post-summer months; therefore, parasite burden also reached maximum levels in summer and autumn. These findings of high prevalence during summer seem to agree closely with the agro climatic conditions of the Kashmir valley. The valley has a temperate climate marked by well-defined seasonality, consisting of four different seasons with wide variations in temperature and other weather conditions that influence the occurrence of parasitic infection in fishes.

Table 1. Season wise prevalence of helminth infection in fishes

Season	Fish Species	No. examined	Positive (%)	P- Value
Spring	<i>S. niger</i>	45	22 (48.88)	0.043
	<i>S. labiatus</i>	30	16(53.33)	
	<i>C. c. communis</i>	41	12(29.26)	
	<i>C. c. spicularis</i>	54	18(33.33)	
Summer	<i>S. niger</i>	27	16(59.25)	0.027
	<i>S. labiatus</i>	36	20(55.55)	
	<i>C. c. communis</i>	36	22(61.11)	
	<i>C. c. spicularis</i>	18	10(55.55)	
Autumn	<i>S. niger</i>	32	20(55.55)	0.037
	<i>S. labiatus</i>	22	10(45.45)	
	<i>C. c. communis</i>	22	6(27.27)	
	<i>C. c. spicularis</i>	27	7(25.92)	
Winter	<i>S. niger</i>	22	7(31.81)	0.028
	<i>S. labiatus</i>	20	5(25.00)	
	<i>C. c. communis</i>	18	3(16.66)	
	<i>C. c. spicularis</i>	36	3(8.33)	
Total		486	197(40.53)	

P < 0.005

DISCUSSION

This pattern of infection is supported by various researchers like Chishty and Peerzada (1998), who while working on seasonal occurrence of acanthocephalan infection in fishes of Wular Lake observed that the infection was higher in summer and low from winter in all fish host. Jha *et al.* (1992) reported that acanthocephalan *Acanthosentis dalti* showed prevalence of 11.1-76% during different months and having highest incidence in the month of May. Frequency of infection and maturation was highest during the may and june as reported by Amin (1975) and Ashley and Nicol (1989) who showed highest prevalence in spring (April-June) and autumn (September-November). Infection patterns of acanthocephalan were greatly influenced by feeding behaviour, temperature, fish species, availability of intermediate host and type of water body. It was seen that overall prevalence rate of *Pomphorhynchus* was low which is in accordance to the studies done by Spall and Summer felt (1969) and Trejo (1994) who showed 0.7% infection of acanthocephalan parasites. The low prevalence might be due to low consumption of intermediate hosts. Seasonal variation in incidence of helminth parasitism in fishes was probably influenced by the annual life cycle of the parasites. In conclusion, these findings needed to be taken into consideration while designing the control strategies for these helminth infections of fishes living under the agro-climatic conditions in the Kashmir valley, as well as similar climatic zones of other parts of world.

REFERENCES

Amin, O. M. 1975. Host and seasonal associations of *Acanthocephalus parjsidei* (Acanthocephala: Echinorhynchidae) in Wisconsin fishes. *J. Parasitol.*, 61(2): 318-329.

- Amundsen, P.A., Knudsen, R., Kuris, A.M. and Kristoffersen, R. 2003. Seasonal and ontogenetic dynamics in trophic transmission of parasites. *Oikos*, 102: 285-293
- Chishty, M. Z. and Peerzada, M. Y. 1998. Host and seasonal occurrence of acanthocephalan in fishes of Wular Lake. *Oriental Science*, 3(1): 31-38.
- Hine, P.M. and Kennedy C.R. 1974. The population biology of the acanthocephala *Pomphorhynchus laevis* (Muller) in the river Avon. *Journal of Fish Biology* 6 665- 679.
- Hussain . A, Khan A.R and Fayaz A. 2003. Future trends in the decline of endemic fish of Kashmir. *Proceedings on Interdisciplinary Approaches in Zoological research*, Kashmir University.
- Jan, N.A and Das, S.M. 1970. Quantitative and qualitative studies on the food of eight fishes of Kashmir Valley. *Ichthyology* 10:21-27.
- Jha, A. N., Sinha, P. and Mishra, T. N. 1992. Seasonal occurrence of helminth parasites in fishes of Sikandarpur reservoir, Muzaffarpur (Bihar). *Indian Journal of Helminthology*, 44(1): 1-8.
- Khanum, H. and Parveen, S. 1997. Organal distribution and seasonal prevalence of endoparasites in *Macrogathus aculeatus* Smith and *Macrogathus armatus* Day. *Bangladesh J. Zool.* 25(1): 15-21.
- Majidah, R. and Khan, A.R. 1998, Seasonal population dynamics of some helminth parasites of fishes in Wular Lake, Kashmir. *Parasitic diseases new horizons Proceedings of the Eighth National Congress of Veterinary Parasitology and National Symposium on Molecular Parasitology*, 34-40.
- Rukhsana, A., Chishty, M.Z., Fayaz, A., Rehana., Bilal, A.B. and Firdousa, A. 2008. Comparative Studies on Helminth Parasites of Declined Fish *Schizothorax esocinus* from two Water Bodies. *4th JK Science Congress* 194.
- Sedar, D.; Mcemal.O. and Yahya. T . 2010. Metazoan parasitological research on three species of fishes collected from canakakkale province north western turkey. *North Western Journal of Zoology*, 6(1): 25-35.
- Tedla, S. and Fernando, C.H. 1969. Observation on the seasonal changes of the parasite fauna of yellow Perch (*Perca flavescens*) from the Bay of Quinte, lake Ontario. *Journal of the Fisheries Research Board of Canada* 26: 866-843.
- Vincent, G. A. and Font, W. F. 2003. Seasonal and yearly population Dynamics of two exotic helminthes, *Camallanus cotti* (Nematoda) and *Bothriocephalus acheilognathi* (Cestoda), parasitizing exotic fishes in Waianu Stream, Hawaii. *The Journal of Parasitology*, 89(4): 756-760.
