



## RESEARCH ARTICLE

### SYMPTOMATOLOGY AND PATHOGENICITY CHARACTERISTICS OF *FUSARIUM OXYSPORUM* F. SP. *CICERI* IN CHICKPEA SEEDS

<sup>1</sup>Vikram, D. Singh and <sup>2</sup>Gangwar, R. K.

<sup>1</sup>Krishi Vigyan Kendra, Ambala (Haryana)

<sup>2</sup>Krishi Vigyan Kendra, Chomu, Jaipur (Rajasthan)

#### ARTICLE INFO

##### Article History:

Received 23<sup>rd</sup> March, 2017

Received in revised form

03<sup>rd</sup> April, 2017

Accepted 18<sup>th</sup> May, 2017

Published online 30<sup>th</sup> June, 2017

##### Key words:

Chickpea, *Fusarium oxysporum* f. sp. *Ciceri*, Pathogenicity.

#### ABSTRACT

*Fusarium* wilt is a serious disease of chickpea in India and other 32 countries. It is soil borne disease. Observation of symptomatology was record. This pathogen was isolated on culture media and natural host, purified and its pathogenicity was proven in pot culture and on potato-dextrose-agar. Pathogenicity of the fungus carried out on a chickpea variety Radhey, which exhibited wilting after 25 days of inoculation. Out of 75 seeds, 68 germinated in infected condition. Symptom appearances were in 20 plants at seedling stage, 14 plants in adult stage, 34 not show any symptom of wilting. Seeds harvested from wilted plants were lighter and duller than those from healthy plants (Haware and Nene, 1980). Further, on the basis of morphological, cultural characteristics of the pathogen and symptomatology, we have confirmed that fungal pathogen, as *Fusarium oxysporum* f. sp. *ciceri*.

Copyright©2017, Vikram, D. Singh and Gangwar. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Vikram, D. Singh and Gangwar, R. K. 2017. "Symptomatology and pathogenicity characteristics of *Fusarium oxysporum* F. sp. *Ciceri* in chickpea seeds", *International Journal of Current Research*, 9, (06), 52435-52437.

#### INTRODUCTION

Pulse crops play an important role in Indian agriculture, besides being rich in proteins; they sustain the productivity of the cropping system. Gram (Chana) (*Cicer aritimum* L.) belongs to the sub family *Papilionaceae* of the family *Leguminaceae*. Chickpea is subjected to attack by a number of fungal, viral, bacterial and nematode diseases. The chickpea wilt caused by *Fusarium oxysporum* f. sp. *ciceri* was reported to be widely distributed in near about 32 countries of the world and at national scenario; six fungal diseases have been reported to be important and causing considerable damage to the crop (Haware *et al.*, 1986a; Nene *et al.*, 1996), causing 10 - 90% annual losses (Singh and Dahiya, 1973; Jalali and Chand, 1992). This pathogen is a facultative saprophyte which can survive in soil up to six years in the absence of susceptible host (Haware *et al.*, 1986). These investigations were undertaken to find out characteristics symptomatology of chickpea wilt complex incidence on field, collection, isolation, purification and pathogenicity of wilt pathogen.

#### MATERIALS AND METHODS

**Symptomatology:** Roots, collar, bark and internal tissues and foliage of diseased part of chickpea plants showing

\*Corresponding author: Gangwar, R. K.  
Krishi Vigyan Kendra, Chomu, Jaipur (Rajasthan)

characteristic symptoms of wilt disease were collected from chickpea field of Plant Pathology section of C. S. A. University of Agriculture and Technology Kanpur at the last week of November, 2002 and March, 2003, for the isolation of *F. oxysporum* f.sp. *ciceri*. Observation of disease symptoms were recorded from start the disease appearance to crop maturity.

#### Isolation, purification, identification and pathogenicity

##### Isolation, purification of pathogen

Isolation of the pathogen wilt affected gram roots, washed in tap water to remove dust particles. The roots were cut into small piece, sterilized with 0.1 per cent mercuric chloride solution for one minute three times. Later these root pieces were transferred in between two folds of sterilized blotter papers to remove the excess of water. The media used was potato dextrose agar. These agar media were prepared by following standard laboratory procedure given by Twite (1969), sterilized by autoclaving, poured into the sterile Petri plates, (ten plates of each medium) and allowed to cool down and solidify. Dried root pieces were finally transferred on two per cent Potato Dextrose Agar (PDA) medium in Petri plates with the help of a sterilized pointed inoculating needle or forceps in the inoculation chamber. These Petri plates were incubated at 28-30<sup>o</sup>C for the growth of the pathogen. When mycelia growth was visible the growing mycelium in different

**Table 1. Effect of wilt pathogen on the seed, seedling and plant of chickpea variety Radhey**

Treatments	No. of seeds germinated	No. of seedlings	No. of plants wilted		
			Seedling stage	Adult stage	Total wilting
Partially sterilized soil inoculated with <i>F. oxysporum</i> f. sp. <i>ciceri</i>	75	68	20	14	34
Partially sterilized soil not inoculated with the fungus (control)	85	85	0	0	0

Petri plates was transferred and purified by repeated transfer of growing hyphae tips to the another sterilized culture tubes and petriplates containing having 2 per cent PDA medium. *F. oxysporum* was purified by single spore method and was identified with the help of relevant literature of Synder and Hausen (1940).

### Identification of the pathogen

The fungus was grown on Potato dextrose agar medium. The measurements of different morphological structures were done under the microscope. Observations on the following morphological characters of the fungus were noted.

- (i) Mycelial characteristics: Colour, branching pattern and width of the hyphae etc., will be studied under microscope for identification.
- (ii) Sporodochia: Its presence, colour, shape, size and arrangement were recorded under the compound microscope.
- (iii) Microconidia: Number, branching pattern, size, shape etc., will be observed under microscope for specific identification.
- (iv) Macroconidia: These will also studied under microscope for number, colour, septation, size and arrangement for specific identification.
- (v) Chlamydospores: Studies on the position, arrangement, shape, and morphology of the Chlamydospores will be done for better understanding about the pathogen.

### Pathogenicity test

Pathogenicity of the fungus was carried out in pot culture by using chickpea variety Radhey which exhibited wilting after 25 days of inoculation. The initial symptoms produced were light yellow and drooping of leaves and final wilting of host. Symptoms of wilting produced by the artificially inoculated and diseased plants were identical and confirmed with those symptoms observed on naturally infected and wilted chickpea plants in the field.

## RESULTS AND DISCUSSION

### Symptomatology

Symptoms of wilt were seen usually in patches of dead seedlings or adult plants seen usually in patches. The disease can affect the crop at any stage. Sudden death, leaves also turn yellow and drop off prematurely, collar region of the wilted plants necrosis and discoloration of the external tissues can be seen. Roots become weak due to disease and when a diseased plant was pulled out, most of the lateral roots remain in the soil. Transverse section of the basal stem or the root reveals masses of fungal hyphae in the vascular bundles and discolorations of vascular cells. The symptoms of chickpea wilt observed were similar to those recorded earlier by Narasimhan (1929), Westerlund et al. (1974), Haware et al. (1986b) and Frisullo et al. (1989).

### Identification of the pathogen

Expansion pathogen on potato-dextrose-agar at 25°C is delicate, white, cottony becoming felted and wrinkled; Mycelium of fungus was profusely bunched, creeping, hyaline, cylindrical, septate and measuring 3.20 - 4.6 µm in width. Colonies of fungus were produces white to light orange sporodochia on potato-dextrose-agar media. Sporodochia was completely covered with arial mycelium, colonies were white initially, radiating in early stage of growth with wine-red pigmentation of medium in later stages which was clearly visible from the bottom side of the plate. Microconidia were hyaline, single celled, oval to cylindrical, straight to slightly curve and measuring 2.5 - 3.5 X 5 - 11 µm in size while macroconidia were sparse and produced on branched macroconidiophores. They were fusoid with pointed ends, hyaline septate (3-4 septate) and 3.0-4.5 X 20-55 µm in size. Chlamydospores were usually intercalary, swell form and one produced singly or in pairs on the hyphae and 7.0 -8 X 3.5 - 5.0 µm. They were globose to sub-globose, thick walled and smooth surfaced. The cultural studies results obtained are in agreement with the findings of Chauhan (1962), Prasad and Patel (1964), Godage (1979) and Kewate (1986).

### Pathogenicity test

In order to ascertain the pathogenic ability of the isolates of *Fusarium oxysporum* f.sp. *ciceri*, the pathogenicity test was conducted on chickpea variety Radhey. The results on pathogenicity test are given in Table-1.

The results presented in table-1 showed that the fungus had adverse effect on the seed germination and seedling emergence. The seed germination was reduced by 9.33%. Out of 75 germinated seeds, 68 seedlings were emerged in pots containing pathogen mixed partially sterilized soil. The reduction in germination and seeding emergence was due do rotting of seeds and pre-emergence rotting of germinated seeds. The initial symptoms of wilting start appearing after 15 days of sowing in the seedlings as drying of seedling from the tip and 20 plants seedlings got infected within two months where as the wilting recorded at the time of adult stage was 14 plants only. The fine roots of the wilted plants were mainly affected and bear dark blacks streak beneath of their bark. Such roots were brittle and easily broken when touched. The brown colour mycelium of the pathogen choked the xylem of primary roots, which resulted in to the drying of whole seedling and plant. The fungus was re-isolated from the diseased tissue and was found same as after comparing with the original one and thus proving the Koch's postulate (1882). Pathogenic variability of *Fusarium oxysporum* f. sp. *Ciceri* has been well documented (Haware and Nene, 1982; Gupta et al., 1986; Honnareddy and Dubey, 2006; Rahman et al., 1998). Different levels of virulence in the populations of *F. oxysporum* f.sp. *lentis* have also been reported by Belabid et al. (2004) and they also did not find any correlation between geographical origin and virulence of the isolates.

## REFERENCES

- Belabid, L., Baum, M., Fortes, Z., Bouznad, Z., Eujayl, I. 2004. Pathogenic and genetic characterization of Algerian isolates of *Fusarium oxysporum* f. sp. *lentis* by RAPD and AFLP analysis. *Afr. J. Biotechnol.*, 3: 25-31.
- Chauhan, S.K., 1962. Physiologic variations in *Fusarium arthoceras* App. and Wr. var. *ciceri* Padwick, causing wilt of gram (*Cicer arietinum* L.) *Proc.Nat. Acad. Sci.*, 32: 78-84.
- Frisullo, S.F., Cioccarese, M., Amenduni, Zamani, H.R. 1989. Wilt of chickpea (*Cicer arietinum* L.) caused by *Fusarium oxysporum* f. sp. *ciceri* in Southern Italy. *Difesa delle Piante*, 12(1-2): 181-185.
- Godage, T.U., 1979. Studies on the wilt of gram (*Cicer arietinum* L.) caused by *Fusarium oxysporum* f. sp. *ciceri* (Padwick) Snyder and HanSen. M.Sc. (Agri.) Thesis submitted to M.P.K.V., Rahuri.
- Gupta, O.M., Khare, M.N., Kotasthane, S.R. 1986. Variability among six isolates of *Fusarium oxysporum* f.sp. *ciceri* causing vascular wilt of chickpea. *Indian Phytopathol.* 39: 279-281.
- Haware, M.P. and Nene, Y.L. 1982. Races of *Fusarium oxysporum* f.sp. *ciceri*. *Plant Dis.*, 66: 809-810.
- Haware, M.P., Nene, Y.L. and Natrajan, M. 1986. Survival of *Fusarium oxysporum* f. sp. *ciceri* in soil in the absence of chickpea. National seminar on management of soil-borne diseases of crop plants. TNAU, Coimbatore, 8-10 Jan. 1986. P.1 (Abstr.).
- Honnareddy, N. and Dubey, S.C. 2006. Pathogenic and molecular characterization of Indian isolates of *Fusarium oxysporum* f.sp. *ciceris* causing chickpea wilt. *Curr. Sci.*, 91: 661-666
- Jalali, B.L. and Chand, H. 1992. Chickpea wilt. In: Singh US, Mukhopadhyay AN, Kumar J, Chaube HS (eds) *Plant Diseases of International Importance. Vol. 1: Diseases of Cereals and Pulses* Prentice Hall, Englewood Cliffs, NJ, pp. 429-444.
- Kewate, S.B. 1986. Studies on wilt of gram (*Cicer arietinum* L.) caused by *Fusarium oxysporum* f. sp. *ciceri*. M.Sc. (Agri.) Thesis. Submitted to M.P.K.V., Rahuri.
- Koch. R. 1882. *Liber die Milzbrand imp. fungus, lime, Emtgegnung auf. den. von. Pasteur in Genf. gehatene Vortrag. T. Fischer. Kassel.*
- Narasimhan, R. 1929. A preliminary note on the *Fusarium* parasitic on Bengal gram (*Cicer arietinum* L). Madras Agri. Dept. Year Book, pp.5-11.
- Nene, Y.L., Haware, M.P. and Reddy, M.V., 1981. Chickpea diseases resistance screening techniques. *Info. Bull.*, 10: 1-4.
- Prasad, N. and Patel, P.N., 1964. Studies on *Fusarium* wilt of Cumin. *Plant Dis. Repr.*, 47: 528-531.
- Rahman, M.L., Haware, M.P., Mian, I.H. and Akananda, A.M. 1998. Races of *Fusarium oxysporum* f.sp. *ciceri* causing chickpea wilt in India. *Bangladesh J. Plant Pathol.*, 14: 234-237.
- Singh, K.B. and Dahiya, B.S., 1973. Breeding for wilt resistance in chickpea. In *Symposium on Problem and Breeding for Wilt Resistance in Bengalgram*, IARI, New Delhi, September 1973. pp. 13-14.
- Snyder, W.C. and Hanson, H.N., 1940. The species concept in *Fusarium*, *Am. J. Bot.*, 27: 64-67.
- Twite, J. 1969. *Plant Pathological Methods. Fungi and Bacteria.* Burgess Publishing Co., pp. 12-80.
- Westerlund, F.V., Jr, Campbell. R.N. and Kimble, K.A. 1974. Fungal root rots and wilt of chickpea in California. *Phytopathology*, 64(4): 432-436.

\*\*\*\*\*