



ISSN: 0975-833X

RESEARCH ARTICLE

ANTIDIABETIC ACTIVITY OF *IPOMEA SEPIARIA* AGAINST STREPTOZOTOCIN (STZ) INDUCED DIABETIC RATS

¹Bamini, B., ²Mani, P., ^{*3}Ganesan, S. and ⁴Natarajan, S.

¹Department of Zoology, Pioneer Kumarasamy College, Nagercoil, Kanyakumari – 629 003, India

²Department of Biotechnology, TUK Arts College, Karanthai, Thanjavur – 613 002, Tamilnadu, India

³Department of Zoology & Biotechnology, A.V.V.M. Sri Pushpam College (Autonomous), Poondi, Thanjavur – 613 503, India

⁴Department of Plant Biology and Plant Biotechnology (Gill Research Institute), Guru Nanak College, Chennai - 600 042, Tamilnadu, India

ARTICLE INFO

Article History:

Received 14th February, 2014

Received in revised form

06th March, 2014

Accepted 19th April, 2014

Published online 21st May, 2014

Key words:

Anti-diabetic,
Hypoglycaemic,
Hypo lipidemic,
Ipomea sepiaria.

ABSTRACT

Diabetes mellitus is a metabolic disorder which results due to chronic hyperglycemia associated with the imbalance in carbohydrate, fat and protein metabolism. Presently available several drugs reduce the hyperglycemia in diabetes mellitus; unfortunately these drugs have side effects. While, herbal drugs are mostly out of toxic or side effect than the chemical drug. Hence, the aim of the present study was designed to compare the possible therapeutic effects of *Ipomea sepiaria* leaves extracts against streptozotocin (stz) induced diabetic rats. The ethanolic extract of *I. sepiaria* was administered orally in an aqueous solution at a dose of 500mg/kg body wt. to diabetic rats. Serum glucose level and lipid profile were estimated after administration of the extracts. Applied doses did not cause any acute toxicity or behavioural changes. The blood glucose levels was significantly ($P < 0.001$) reduced when compared to the streptozotocin (STZ) induced diabetic rats. The lipid profile such as Total Cholesterol, Total Glycerides, Low Density Lipoproteins and Very Low Density Lipoproteins levels were significantly decreased in *I. sepiaria* treated diabetic animals. In contrast, High Density Lipoproteins levels were increased when compared to the Diabetic control rats. In conclusion, *I. sepiaria* leaves produced a significant hypoglycemic effect and also hypolipidemic activity at dose level of 500 mg/kg.

Copyright © 2014 Bamini et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Diabetes mellitus is an endocrine metabolic disorder characterized by hyperglycemia, altered lipids, carbohydrates, proteins metabolism and it increases risk of cardiovascular diseases complications (Davis, 2006). The two forms of diabetes, type 1 and 2, differ in their basic mechanisms of development and in physiologic characteristics such as associations with obesity, age, and insulin. But, both types of the diabetes share the common characteristics of hyperglycemia, microvascular and macrovascular complications. Moreover, the alterations of lipoproteins metabolism are involved to the pathogenesis of the cardiovascular disease in both forms of diabetes in a similar way (Howard, 1987). Diabetes has a considerable impact on the health, life style, life expectancy of patients and its related complications are major healthcare problems. Currently, diabetes is controlled by handful of available drugs such as oral hypoglycemic agents and insulin, but they have their own limitations. Traditionally, many herbal medicines and

medicinal plants have been used for the treatment of diabetes as an alternative medicine (Mukherjee *et al.*, 2006). Presence of various phytoconstituents in medicinal plants is thought to act on a different series of targets by multiple modes and mechanisms. Hence, plants have the potential to impart therapeutic effect in complicated disorders like diabetes and its complications (Tiwari and Rao, 2002). Screening of medicinal plants is one of the alternative and valid approaches in the drug development process because they contain diverse phytoconstituents which may give new drug leads and may be effective and safe in diabetes. In India, traditionally numbers of plants are used to manage the diabetic conditions and their active principles were isolated but few plants have been scientifically studied. Therefore, the present study was carried out to evaluate the antidiabetic activity of *I. sepiaria* in STZ induced diabetes and to probe into the mechanism of its antidiabetic property.

MATERIALS AND METHODS

Animals

Male albino rats of Wistar strain approximately weighing 160-180g were used in this study. They were healthy animals

*Corresponding author: Ganesan, S.

Department of Zoology & Biotechnology, A.V.V.M. Sri Pushpam College (Autonomous), Poondi, Thanjavur – 613 503, India.

purchased from the Indian Institute of Science, Bangalore. The animals were housed in spacious polypropylene cages bedded with rice husk. The animal room was well ventilated and maintained under standard experimental conditions (Temperature $27 \pm 2^\circ$ C and 12 hour light/dark cycle) throughout the experimental period. All the animals were fed with standard pellet diet and water were provided *ad libitum*. They were acclimatized to the environment for one week prior to experimental use. The animal feed composition is crude protein (22.3%), crude oil (4.01%), crude fibre (4.02%), Ash (8.02%) and sand silica (1.02%).

Chemicals

Streptozotocin (STZ), Ethylene Diamine Tetra Acetic Acid (EDTA), Glibenclamide (Prudence Pharma Chem, India), Chloroform were purchased for Sigma chemical company, Mumbai All other chemicals and reagents used in this study was of analytical grade with high purity and were obtained from Glaxo laboratories and Sisco Research laboratories, Mumbai, India.

Preparation of plant extract

The leaves of *I. sepiaria* were collected from in and around Thanjavur, Tamil Nadu, India and identified with help of the standard manuals such as "The Flora of the presidency of Madras" (Gamble, 1967) The collected leaves of *I. sepiaria* were cut into small pieces and shade dried at room temperature and makes a fine powder using grinder mixture. The powder material of *I. sepiaria* leaves were macerated with 70% ethanol at room temperature for 3 days. After 3 days, the supernatant was transferred into china dish. The supernatant was completely removed by keeping the china dish over a boiling water bath at 45 C. A semi solid extract was obtained after complete elimination of alcohol. The obtained residue was kept in the refrigerator for further use. The extract was made up to a known volume in distilled water just before oral administration.

Streptozotocin (STZ) Induced Diabetic rats

The animals were divided into four groups of six animals each. Diabetes was induced in all groups except normal control following overnight fasting (deprived of food for 16h allowed free access to water) by a single intraperitoneal injection of 65mg/kg of streptozotocin (STZ) dissolved in a freshly prepared 0.1M citrate buffer (pH4.5) (Liu et al., 2008). The animals of normal control (Group I) were injected with saline alone. Diabetes was confirmed 72 h after induction by measurement of tail vein blood glucose levels by glucose oxidase-peroxidase method using strips. Group II served as diabetogenic rats (Control). Group III rats treated with *I. sepiaria* at a dose of 500mg/kg was orally given once a day for 15 days after hyperglycemia was confirmed. Group IV rats treated with Glibenclamide as a standard at dose of 0.25mg/kg (Arulmozhi et al., 2010). After complete the experimental period, the animals were killed cervical dislocation after an overnight fasting. The blood sample was collected. The blood was allowed to clot by standing at room temperature for 30 minutes and then refrigerated for another 30 minute. The

resultant clear part was centrifuged at 3000 rpm for 10minutes and then the serum (supernatant) was isolated and stored at refrigerated until required for analysis.

Biochemical estimations

Serum glucose was estimated by the oxidase method (Trinder, 1969). The total cholesterol was estimated by the method of Allain et al., (1974). Triglyceride was estimated by the method of Werner et al., (1981). HDL cholesterol was separated by adding phosphotungsti magnesium chloride to the fresh samples to precipitate other lipoproteins and the HDL cholesterol was estimated by the method of Allain et al., (1974). The concentration of LDL cholesterol was calculated by using the Friedwald formula (1972) and VLDL cholesterol was calculated by dividing the triglycerides value (in mg/dl) by 5. Hemoglobin estimated by the method of Dacie and Lewis, (1968).

Statistical Analysis

Values were expressed as mean \pm standard deviation for six rats in the each group and statistical significant differences between mean values were determined by one way analysis of variance (ANOVA) followed by the Tukey's test for post-hoc multiple comparison tests. Statistical Package for Social Studies (SPSS) 9.0 version was used and $p < 0.001$ was considered to be significant.

RESULTS

The ethanolic extract of *I. sepiaria* was administered orally in an aqueous solution at a dose of 500mg/kg body wt. to diabetic rats to assess the synergetic impact of the plant extracts. The plant extracts were fed with normal and diabetes induced rats. The blood glucose levels was significantly ($P < 0.001$) reduced when compared to the specific diabetic control animals (Table 1).

The lipid profile such as TC, TG, LDL and VLDL levels were significantly increased in diabetic control animals (DC) where as HDL levels were decreased when compared to the control rats. The plant extracts were administered orally at a dose of 500mg/kg body wt., to diabetic rats significant ($P < 0.001$) depletion in the total cholesterol, TG, LDL, and VLDL levels and increment of HDL levels were recorded in the diabetic animals (Table 1 and 2). The depleted high density lipoprotein (HDL) in the diabetic rats, increased significantly ($P < 0.001$) after the administration of the plant extract. The plant extract possesses significant antidiabetic activity and close proximity to standard.

DISCUSSION

Streptozotocin (STZ) (2-deoxy-2-(3-methyl-3-nitrosuureido)-D-glucopyranose) is commonly used for experimental induction of type-I diabetes mellitus, which causes selective pancreatic islet β -cell cytotoxicity mediated through the release of nitric oxide (NO). This results in rapid reduction in pancreatic islet pyridine nucleotide concentration and subsequent β -cell necrosis. The action of STZ on mitochondria generates SOD anions, which leads to diabetic complications

Table 1. shows the effect of *I. sepiaria* on glucose, Hb, cholesterol and triglycerides in experimental rats

Treatment Groups	Glucose (mg/dl)	Hb (g/dl)	Total cholesterol (mg/dl)	Triglycerides (mg/dl)
Group I	101.69 ± 5.08	13.76 ± 0.68	90.96 ± 4.54	114.28 ± 5.71
Group II	206.32 ± 10.31 ^a	9.03 ± 0.45 ^a	236.36 ± 11.81 ^a	188.57 ± 9.42 ^a
Group III	96.61 ± 4.83 ^b	15.49 ± 0.77 ^b	76.36 ± 3.81 ^b	82.85 ± 4.14 ^b
Group IV (S)	98.30 ± 4.91 ^b	13.35 ± 0.66 ^b	82.74 ± 4.13 ^b	87.14 ± 4.35 ^b

Values were expressed as mean ± SD for six rats each.

^a Compared with group I (p<0.001) ^b Compared with group II (p<0.001)

Table 2. shows the effect of *I. sepiaria* on HDL, VLDL, and LDL- cholesterol in Experimental Rats

Treatment Groups	HDL cholesterol (mg/dl)	VLDL cholesterol (mg/dl)	LDL cholesterol (mg/dl)
Group I	24.57 ± 1.22	28.85 ± 1.44	37.54 ± 1.87
Group II	18.03 ± 0.98 ^a	37.71 ± 1.88 a	180.26 ± 8.18 a
Group III	27.11 ± 1.35 ^b	16.57 ± 0.62 ^b	37.76 ± 0.08 ^b
Group IV (S)	25.86 ± 1.29 ^b	17.42 ± 0.57 ^b	43.86 ± 2.19 ^b

Values were expressed as mean ± SD for six rats each.

^a Compared with group I (p<0.001) ^b Compared with group II (p<0.001)

(Papaccio *et al.*, 2000). Based on the above perspectives, in the present study, the antidiabetic activity has been assessed in rats made diabetic by STZ. Sulfonyl ureas such as glibenclamide are often used as a standard antidiabetic drug in STZ-induced diabetes to compare the efficacy of variety of antihyperglycemic compounds (Anderson *et al.*, 1974). In diabetes the increased blood sugar levels might be due to either insulin resistance of the body cells or decreased secretion of insulin from beta cells manifest in the decreased serum insulin levels (Mohammad Ali, *et al.*, 2004). Administration of G1, G2, G3, G4, G5 and G6 to diabetic rats restored the levels of glucose. Present finding is in agreement with Subramaniam *et al.*, (2012) studies. Diabetes affects both glucose and lipid metabolism (Sperling *et al.*, 2000). The insulin deficiency depletes the activity level of lipoprotein lipase, thus leading to deranged lipoprotein metabolism during diabetes (Ranganathan *et al.*, 2000).

The lipoprotein levels in the STZ induced diabetic rats of the present study reveal a significant alter in lipoprotein metabolism. The serum total cholesterol content increased significantly in diabetic animals. The elevated hypertriglyceridemia was increased in the synthesis of triglyceride rich lipoprotein particles (very low density lipoprotein, VLDL) in liver diminished catabolism in diabetic rats (Ginsberg, 1991). Since insulin has a potent inhibitory effect on lipolysis in adipocytes, insulin deficiency is associated with excess lipolysis and increased influx of free fatty acids to the liver (Coppack, 1994, Ohno, 2000) The increased levels of low-density lipoprotein (LDL) and very low density lipoprotein (VLDL) in the diabetic animals might be due to over production of LDL and VLDL by the liver due to the stimulation of hepatic triglyceride synthesis as a result of free fatty acid influx (Coppack, 1994). The high density lipoprotein (HDL) was significantly reduced in the diabetic rats which indicate a positive risk factor for atherosclerosis (Bopanna, 1997). Supplementations of *I. sepiaria* to diabetic rats restored the lipid profile. Our results concord with the earlier work done by Kesari *et al.*, (2006), where it has been reported that lipid profile level in the plasma is restored with the treatment of *Aegle marmelos* seed extract in diabetic rats.

The blood glucose level of *I. sepiaria* extracts fed animal was significantly ($P<0.001$) reduced. The levels of serum TC, TG, LDL, and VLDL were found to be significantly reduced in the plant extracts treated diabetic animals. This might be due to the reduced hepatic triglyceride synthesis and or reduced lipolysis that might be due to the increase in serum insulin levels in the plant extract treated rats. The HDL increased significantly in the plant extract treated rats indicating a reversed atherogenic risk. In uncontrolled or poorly controlled diabetes there is an increased glycosylation of a number of proteins including haemoglobin and -crystalline of lens (Alberti and Press, 1982). Glycosylated haemoglobin (HbA1) was found to increase in patients with diabetes mellitus to approximately 16% (Koenig *et al.*, 1976) and the amount of increase is directly proportional to the blood glucose level (Jackson *et al.*, 1979). During diabetes the excess glucose present in blood reacts with haemoglobin to form glycosylated haemoglobin. Therefore, the total haemoglobin level is decreased in STZ diabetic rats. So the total haemoglobin level is lowered in alloxan diabetic rats (Sheela and Augusti, 1992). Administration of *I. sepiaria* reversed the total haemoglobin levels in alloxan diabetic rats. The present study suggests that the *I. sepiaria* extracts had synergetic hypoglycemic effect revealed by decreased serum lipid levels, restored haemoglobin and therefore attribute to therapeutic value of the plant extracts of *I. sepiaria* to combat the diabetic condition in rats.

REFERENCES

- Alberti KGMM & Press CM 1982. In: Keen, H., Jarre, J. ZEds. The biochemistry and the complications of Diabetes. Edward Arnold Publishers, pp. 231_270.
- Allain CC, Poon LS, Chan CSG, Richmond W & Fu PC 1974. Enzymatic determination of total serum cholesterol. Clinical Chemistry 20: pp 470-475.
- Anderson T, Schein PS, McMenamin MG & Cooney DA 1974. Streptozotocin diabetes correlation with extent of depression of pancreatic islet nicotinamide adenine dinucleotide. *Journal of Clinical Investigation*, 54: 672-677.

- Arulmozhi S, Mazumder PM, Lohidasan S, Thakurdesai P 2010. Antidiabetic and antihyperlipidemic activity of leaves of *Alstonia scholaris* Linn. R.Br. *European Journal of Integrative Medicine*. (2)(1) 23-32.
- Bhagavan N.V. 2002. Metabolic homeostasis in: N.V. Bhagavan (Eds) *Medical biochemistry* Harcourt/Academic, Florida, 485-519.
- Bopanna KN, Kannan J, Sushma G, Blaraman R & Rathod SP 1997. Antidiabetic and anti hyperlipidemic effect of Neem, Lipidemic effect of Neem seed kernel powder on alloxan diabetic rabbits. *Ind. J. Pharmacology*, 29: 162-167.
- Coppack SW, Jenson MD & Miles JM 1994. In vivo regulation of lipolysis in human. *J. lipid Res*. 35:177-193.
- Dacie JV & Lewis SM (1968). *Practical Hematology*, 4th edition J and A, Churchill, UK. p37.
- Davis S. 2006. Insulin, oral hypoglycemic agents and the pharmacology of the endocrine pancreas. In: Brunton L, Lazo J, Parker K. (eds.) *Goodman and Gilman's the pharmacological basis of therapeutics*. New York: McGraw Hill, 1613-1646.
- Duran RG, Pascoe LD, Vilar RC & Medina N.R. 2004. Diabetogenic effect of STZ diminishes with loss of Nitric oxide role of ultra violet and carboxy PT10 pharmacology, 71(1)17-24.
- Friedwald's WT, Levy RT & Fredrickson DS 1972. Estimation of low-density lipoprotein cholesterol in plasma, without use of the preparative centrifuge. *Clin. Chem*, 23: 499.
- Gilman AG, Rall TW, Nles AS, Tayer P 1990. *The pharmacological basis of therapeutics*, eighth edn. Pergamen press, Newyork, 1317-1322 and 1463-1495.
- Ginsberg HN 1991. Lipoprotein physiology in nondiabetic and diabetic states. Relationship to atherogenesis. *Diabetes care*, 14: 839-855.
- Harvey J & Paige SM 1998. *The Instat Guide to choosing and interpreting statistical tests: A manual for Graph pad Instat, Version 3*. San Diego, CA USA.
- Howard BV 1987. Lipoprotein metabolism in diabetes mellitus. *J Lipid Res*, 28: 613-628.
- Jackson RL, Hess RL & England JD 1979. Haemoglobin A1C values in children with overt diabetes maintained in varying degree of control. *Diabetes Care*, 2: 391-395.
- Jarvenin YH 1995. Role of insulin resistance in the pathogenesis of NIDDM. *Diabetologia*, 38: 1378-1388.
- Kesari AN, Rajesh KG, Santosh KS, Sandhya D & Geeta W 2006. Hypoglycemic and antihyperglycemic activity of *Aegle marmelos* seed extract in normal and diabetic rats. *Journal of Ethnopharmacology*, 107: 374-379.
- Koenig RJ, Peterson CM, Jones RL, Saudek C, Lehrman M & Cerami A 1976. Correlation of glucose regulation and haemoglobin A1C in diabetes mellitus. *New Engl. J. Med*, 295: 417-420.
- Kurup S & Bhone RR 2000. Combined effect of nicotinamide and streptozotocin on diabetic status in partially pancreatectomized adult BALB/C mice. *Horm Metab Res*, 32(8) 330-334.
- Liu H, Liu X, Lee J, Liu Y, Yang H & Wang G 2008. Insulin therapy restores impaired function and expression of P-glycoprotein in blood-brain barrier of experimental diabetes. *Biochemical Pharmacology*, 75:1649-58.
- Mohammad AE & Razeih Y 2004. Hypoglycemic effect of *Teucrium polium* studies with rat ancreatic islets. *J. Ethnopharmacol*, 95:27-30.
- Mukherjee P, Maiti K, Mukherjee K & Houghton PJ 2006. Leads from Indian medicinal plants with hypoglycemic potentials. *J Ethnopharmacol*, 106: 1-28.
- Ohno T, Horio F, Tanaka S, Terada M, Namikawa T & Kitoh J. 2000. Fatty liver and hyperlipidemia in IDDM of streptozotocin treated shrews. *Life Sci*, 66:125-131.
- Papaccio G, Pisanthi FA, Latronico MY, Ammendola E & Galdieri M. 2000. Multiple low-dose and single high dose treatments with streptozotocin do not generate nitricoxide. *Journal of Cellular Biochem*, 77:82-91.
- Ranganathan G, Li C & Kern PA 2000. The translational regulation of lipoprotein lipase in diabetic rats involves the 3'-untranslated region of lipoprotein lipase mRNA. *J.Biol.Chem*, 275:40986-40991.
- Sperling MA & Saunders PA 2000. *Diabetes mellitus in: R.E. Behrman R.M, Kliegman H.B, Jenson (Eds) Nelson text book of pediatrics*, 1767-1791.
- Subramaniam R, Aiyalu R & Manisenthilkumar KT 2012. Investigation of hypoglycemic, hypolipidemic and antioxidant activities of aqueous extract of *Terminalia paniculata* bark in diabetic rats. *Asian Pacific Journal of Tropical Biomedicine*, 262-268
- Tiwari A & Rao J 2002. Diabetes mellitus and multiple therapeutic approaches of phytochemicals: present status and future prospects. *Curr Sci*, 83: 30-38.
- Tiwari AK & Madhusudana RJ 2002. Diabetes mellitus and multiple therapeutic approaches of phytochemicals: present status and future prospects. *Current Science*, 83: 30-38.
- Trinder P 1969. *Practical Clinical Biochemistry*, Vol x, 5th edit., William Heinemann Medical Books Limited, New York.
- Werner M, Gabrielson DG & Eastman G 1981. Ultramicro determination of serum triglycerides by bioluminescent assay. *Clinical Chemistry*, 27:268-271.
- Yoon JW & Ray UR 1985. Perspectives on the role of viruses in insulin dependent diabetes. *Diabetes care*, 1:39-44.
