



RESEARCH ARTICLE

CLINICAL EVALUATION OF CONNECTIVE TISSUE GRAFT IN THE TREATMENT OF MILLERS CLASS I & II MULTIPLE ADJACENT GINGIVAL RECESSIONS IN THE ESTHETIC ZONE

¹Dr. Syed Saima, ²Dr. Dolly, ³Dr. Aqrib Mushtaq and ⁴Dr. Mumtaz Ali

¹Lecturer, Dept. of Periodontics, Govt. Dental College, Srinagar

^{2,3,4}Post-graduate Scholar, Dept. of Periodontics, Govt. Dental College, Srinagar

ARTICLE INFO

Article History:

Received 24th February, 2019

Received in revised form

29th March, 2019

Accepted 10th April, 2019

Published online 22nd May, 2019

Key Words:

Gestation Length,
Dangi, Parity, Body Score Condition.

ABSTRACT

Background: Multiple approaches have been used to replace lost, damaged or diseased gingival tissues. The connective tissue graft (CTG) procedure is the gold standard method for root coverage. Although multiple sites often need grafting, the palatal mucosa supplies only a limited area of grafting material. To overcome this limitation, expanded mesh graft provides a method whereby a graft can be stretched to cover a large area. The aim of this study was to evaluate the effectiveness and the predictability of expanded mesh CTG (e-MCTG) in the treatment of adjacent multiple gingival recessions. **Materials and Methods:** Sixteen patients aged 20-50 years contributed to 55 sites, each site falling into at least three adjacent Miller's Class 1 or Class 2 gingival recession. The CTG obtained from the palatal mucosa was expanded to cover the recipient bed, which was 1.5 times larger than the graft. Clinical measurements were recorded at baseline and 3 months.

*Corresponding author:

Copyright © 2019, Syed Saima et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Dr. Syed Saima, Dr. Dolly, Dr. Aqrib Mushtaq and Dr. Mumtaz Ali, 2019. "Clinical evaluation of connective tissue graft in the treatment of Millers class I & II multiple adjacent gingival recessions in the esthetic zone", *International Journal of Current Research*, 11, (05), 3900-3903.

INTRODUCTION

Gingival recession is defined as the partial denudation of the root surface due to the apical migration of the gingival margin. (Guiha, 2001) Etiological factors include trauma from tooth brushing, malposition of teeth, ectopic insertion of frenum, and muscle attachments. The major therapeutic goals in mucogingival surgery are a correction of esthetic problems and management of hypersensitivity. Numerous surgical procedures have been described to achieve soft tissue coverage of exposed root surfaces including coronally repositioned flaps, pedicle grafts, free gingival grafts, sub epithelial connective tissue grafts (CTG), and guided tissue generation (GTR) <http://www.contempclindent.org/article.asp?issn=0976-237X;year=2015;volume=6;issue=3;page=321;epage=326;aulast=Shanmugam#ref2> were commonly used procedures. (Cordioli, 2001; Romagna-Genon, 2001; Danesh-Meyer, 2001; Jahnke, 1993; Pini Prato et al., 2000; Casati et al., 2000; da Silva Pereira, 2000; Pini Prato, 1992; Saletta, 2001). The treatment of isolated or multiple buccal recessions with different surgical procedures depends on many factors such as defect size, presence or absence of keratinized tissue adjacent to the defect, and thickness of the gingiva. Since the patients are concerned about their esthetic appearance, every effort should be made to achieve complete root coverage up to the cemento-enamel junction (CEJ).

(Danesh-Meyer, 2001; Zucchelli, 2001; Henderson, 2001). Originally Sullivan and Atkins (Sullivan, 1968) described a technique for coverage of exposed root surfaces using the free gingival autogenous graft. The graft survival over large expanses of avascular root surfaces was unpredictable, and complete root coverage was rarely achieved. Karring et al. (1972) demonstrates that the underlying connective tissue has a direct bearing on the type of epithelium that is superimposed upon it. Edel (1974) showed that a significant increase in the volume of gingiva can be achieved by grafting (Pini Prato, 1992; Sullivan, 1968) gingival connective tissue alone. Langer and Langer (Langer, 1985) described the CTG technique in root coverage on both single and multiple adjacent teeth. The advantage is the dual blood supply from the overlying flap and palatal connective tissue, which maximizes graft survival. It also provides excellent esthetic results. When multiple adjacent teeth with gingival recessions are present in esthetic regions of the mouth, the preferred surgical technique should be such the one, which provides the possibility of achieving maximum root coverage. One of the problems with multiple root coverage grafting is the unavailability of the large blood supply of donor tissue. If connective tissue supply is limited, more than one surgical procedure may be needed. (Zucchelli, 2000; Henderson, 2001). The purpose of the present study was to evaluate the effectiveness and the predictability of expanded

mesh CTG (e-MCTG) procedure for the treatment of multiple adjacent gingival recession.

MATERIALS AND METHODS

The study population 16 patients, (age range 20-55 years mean age 37 years) with either dentin hypersensitivity or esthetic problems caused due to the recession defects were included in the study. A total of 55 sites were treated in 16 patients. Prior to initiation of the study, ethical approval was obtained from institution ethical committee. All the patients agreed to the study protocol, and signed informed consent was obtained prior to inclusion in the study. The inclusion criteria are

- The presence of at least three adjacent Miller's Class I or Class II gingival recession (Miller, 1985) on the buccal/facial aspect with recession depth (RD) of ≥ 2 mm,
- Probing depth (PD) of ≤ 3 mm.
- A minimum width of keratinized gingival (KG) of at least 1 mm. Nine subjects contributes three sites, and seven subjects contributed four sites defects.
- The exclusion criteria are (1) the presence of severe cervical abrasion/root caries,
- The presence of abnormal frenal attachment, (3) Current smokers, (4) Medically compromised patients, (5) Miller's Class III and IV gingival recession. The patients initially completed a plaque control program, so as to achieve a full mouth plaque score (FMPS) $< 25\%$.

Clinical measurements

The following clinical measurements were taken by a single examiner at baseline and 3 months

- RD measured from the Cemento-Enamel Junction (CEJ) to the gingival margin
- Recession width (RW) measured across the buccal surface at the CEJ level
- PD measured from the gingival margin to the bottom of the gingival sulcus
- Width of keratinized tissue (KT) measured from the gingival margin into the mucogingival junction
- Clinical attachment level (CAL) measured from CEJ to the bottom of the gingival sulcus.

All measurements were performed at the mid buccal level using a William's periodontal probe (Hu-Friedy) and rounded to the nearest 0.5 mm

Surgical procedure: All surgical procedures were done by the same operator. Following the induction of local anesthesia (Lignocaine hydrochloride with 1:100,000 adrenaline), an intra-crevicular incision was made through the bottom of the crevice and horizontal incision was placed at the level of CEJ extending 3 mm on either side of the involved tooth including their papilla. Two vertical incisions were placed from the end point of the horizontal incision to the alveolar mucosa to establish a trapezoidal flap. A full thickness flap was elevated to 3-4 mm apical to the bone dehiscence followed by a split thickness flap and all muscle interferences were eliminated in order to facilitate its coronal advancement. The remaining buccal soft tissue of the anatomic interdental papillae was de-

epithelized. The root surface was mechanically instrumented using Gracey curettes followed by conditioning with 1 ml tetracycline hydrochloride solution for 3 min with subsequent rinsing with saline. CTG was harvested from the molar-premolar area of the palate on one side and The donor site was then sutured with 4-0 black silk to ensure primary intention healing. Alternating incisions were then made on each edge of the harvested graft to expand it so that it would cover the recipient bed completely, which was 1.5 times larger than the graft. Subsequently, the graft was positioned at the CEJ with interrupted 5-0 vicryl bioabsorbable sutures. The mucogingival flap was coronally repositioned without tension to cover the e-MCTG with 4-0 silk sutures. The area was re-examined to ascertain that the graft was completely covered by the flap. A periodontal dressing (coe-pak) was placed over the recipient site and removed after a week. All patients were instructed to discontinue tooth brushing in the surgical site for 1-week so as to avoid trauma or pressure at the surgical site. A 0.12% chlorhexidine digluconate mouth rinse was prescribed 2 times daily for 15 days. Analgesics (Ibu Profen and paracetamol tds for 5 days) and antibiotics (amoxycillin 500 mg tds for 5 days) were prescribed. Mechanical tooth cleaning of the treated areas using a soft toothbrush and a careful roll technique was resumed following the removal of periodontal dressing. The patients were recalled for oral prophylaxis after 2, 4 weeks, and every 3 months.

RESULTS

At baseline, mean RD was $2.56 \text{ mm} \pm 0.62 \text{ mm}$, mean RW was $3.44 \text{ mm} \pm 0.44 \text{ mm}$, mean KT was $2.13 \text{ mm} \pm 0.73 \text{ mm}$, mean PD was $1.15 \text{ mm} \pm 0.23 \text{ mm}$, and a mean CAL was $3.71 \text{ mm} \pm 0.61 \text{ mm}$. Three months following surgical intervention, the mean RD reduced from $2.56 \text{ mm} \pm 0.62 \text{ mm}$ to $0.61 \text{ mm} \pm 0.63$. (mean $1.96 \text{ mm} \pm 0.66 \text{ mm}$), the mean root coverage was 86%, the mean RW reduced from $3.44 \text{ mm} \pm 0.44 \text{ mm}$ to $1.52 \text{ mm} \pm 1.40 \text{ mm}$ (mean 1.92 ± 1.34), the mean KT increased from $2.13 \text{ mm} \pm 0.73 \text{ mm}$ to $3.55 \text{ mm} \pm 0.69 \text{ mm}$ (mean 1.42 ± 0.62), the PD from $1.15 \text{ mm} \pm 0.23 \text{ mm}$ to $1.22 \text{ mm} \pm 0.25 \text{ mm}$ (0.07 ± 0.33), and CAL increased from $3.71 \text{ mm} \pm 0.61 \text{ mm}$ to $1.83 \text{ mm} \pm 0.70 \text{ mm}$ (1.88 ± 0.69). On statistical analysis, there was a significant reduction in RD and RW, and KT and clinical attachment gain at 3 months ($P < 0.001$) compared to the

DISCUSSION

Gingival recession involves groups of adjacent teeth and is seldom localized to a single tooth. When multiple recession defects affecting adjacent teeth in esthetic areas of the mouth are present, they should all be treated at the same time to help ensure the best esthetic results. Autogenous CTG have been extensively used for root coverage procedures in teeth and implants (Borghetti, 1994; Bouchard, 1994) Whereas subepithelial CTG was extensively used for one or two adjacent gingival recession defects excellent result with color matching. (Langer, 1985) Harris's study proposed that the use of acellular dermal matrix graft would improve the gingival color, reduce patient morbidity, provide a uniform thickness of material and eliminate the need for multiple surgeries because of unlimited availability (Aichelmann-Reidy, 2001). In the present study, a new approach of the CTG technique was described to cover multiple gingival recession defects. The most common problem for root coverage with CTG procedure is the amount that can be harvested.



Pre-operative View



Incision given on donor site



Harvesting of CTG



Preparation of recipient site



Suturing done



Post-operative View

The rugae area is not suitable for graft material, and an extensive palatal wound will be uncomfortable for the patient. Because of the high rate of complications and a limited amount of palatal mucosa available for grafting, it is advisable to refrain from covering large or multiple defects. Formerly e-MCTG was used for free gingival grafts and was generally applied to increase the width of keratinized tissue without root coverage (Rateitschak, 1989). We modified this technique to cover multiple gingival recession sites in one surgery. E-MCTG provides more graft material since it can be expanded as much as 50% to cover a larger area. This surgical technique resulted in complete root coverage in 44 out of 55 sites. This success rate is similar to those previously reported by Cordioli *et al.*, (2001) 94.68%; Romangno-Genon (2001) 84.84%; Rosetti *et al.*, (2000) 95.6%; and Harris (1994) 97.7%. The free gingival graft is commonly applied for increasing the width of keratinized tissue (Jahnke, 1993; Harris, 2001).

However, it has some limitations and complications such as color match, painful postoperative wound healing, and scar tissue formation in the donor area. It has been reported that using CTG to increase KT has a more rapid, maturation, and less traumatic healing of the graft in the recipient site. (Jahnke, 1993; Rosetti, 2000; Harris, 2001) Similar clinical observations were noticed in our study with the use of the e-MCTG technique. The tissue was tightly bound to the tooth in most cases and resisted probing. Histological studies have demonstrated that the blood supply from the periosteum and overlying flap results in a more rapid re-establishment of circulation (Guiha, 2001; Goldstein, 2001). In the present study, we took care to place the graft with the periosteal side facing the root surface. Recently, an acellular dermal matrix has been shown to be effective in root coverage procedures as a substitute for CTG. Tat *et al.* (2002) and Wei *et al.* (2000) have shown that acellular dermal matrix was not as successful

as the autogenous free graft and connective tissue free graft in increasing the KT, and a histologic report suggested that placing an acellular dermal matrix does not increase KT.

Conclusion

CTG is the gold standard for treatment of gingival recession, but the disadvantages are the inadequate graft availability. The results of the present study demonstrated that the e-MCTG procedure was an effective and predictable treatment modality for the management of multiple adjacent gingival recessions in terms of root coverage in the treatment of multiple adjacent gingival recessions.

REFERENCES

- Aichelmann-Reidy ME., Yukna RA., Evans GH., Nasr HF., Mayer ET. 2001. Clinical evaluation of acellular allograft dermis for the treatment of human gingival recession. *J Periodontol.*, 72:998-1005.
- Borghetti A., Gardella JP. 1990. Thick gingival auto graft for the coverage of gingival recession, clinical evaluation dentistry. *Int J Periodontics Restorative Dent.*, 10:216-29.
- Borghetti A., Louise F. 1994. Controlled clinical evaluation of the subpedicle connective tissue graft for the coverage of gingival recession. *J Periodontol.*, 65:1107-12.
- Bouchard P., Etienne D., Ouhayoun JP., Nilvéus R. 1994. Subepithelial connective tissue grafts in the treatment of gingival recessions. A comparative study of 2 procedures. *J Periodontol.*, 65:929-36.
- Casati MZ., Sallum EA., Caffesse RG., Nociti FH. Jr, Sallum AW., Pereira SL. 2000. Guided tissue regeneration with a bioabsorbable polylactic acid membrane in gingival recessions. A histometric study in dogs. *J Periodontol.*, 71:238-48.
- Cordioli G., Mortarino C., Chierico A., Grusovin MG., Majzoub Z. 2001. Comparison of 2 techniques of subepithelial connective tissue graft in the treatment of gingival recessions. *J Periodontol.*, 72:1470-6.
- da Silva Pereira SL, Sallum AW, Casati MZ, Caffesse RG, Weng D., Nociti FH. Jr, et al. 2000. Comparison of bioabsorbable and non-resorbable membranes in the treatment of dehiscence-type defects. A histomorphometric study in dogs. *J Periodontol.*, 71:1306-14.
- Danesh-Meyer MJ., Wikesjö UM. 2001. Gingival recession defects and guided tissue regeneration: A review. *J Periodontol Res.*, 36:341-54.
- Edel A. 1974. Clinical evaluation of free connective tissue grafts used to increase the width of keratinised gingiva. *J Clin Periodontol.*, 1:185-96.
- Goldstein M., Boyan BD., Cochran DL., Schwartz Z. 2001. Human histology of new attachment after root coverage using subepithelial connective tissue graft. *J Clin Periodontol.*, 28:657-62.
- Guiha R. el Khodeiry S., Mota L., Caffesse R. 2001. Histological evaluation of healing and revascularization of the subepithelial connective tissue graft. *J Periodontol.*, 72:470-8.
- Harris RJ. 1994. The connective tissue with partial thickness double pedicle graft: The results of 100 consecutively-treated defects. *J Periodontol.*, 65:448-61.
- Harris RJ. 2001. Clinical evaluation of 3 techniques to augment keratinized tissue without root coverage. *J Periodontol.*, 72:932-8.
- Henderson RD., Greenwell H., Drisko C., Regennitter FJ., Lamb JW., Mehlbauer MJ. et al., 2001. Predictable multiple site root coverage using an acellular dermal matrix allograft. *J Periodontol.*, 72:571-82. †
- Jahnke PV., Sandifer JB., Gher ME., Gray JL., Richardson AC. 1993. Thick free gingival and connective tissue autografts for root coverage. *J Periodontol.*, 64:315-22. †
- Karring T., Lang NP., Loe H. 1972. The role of gingival connective tissue in determining epithelial differentiation. *J Dent Res.*, 51:1303-4.
- Langer B., Langer L. 1985. Subepithelial connective tissue graft technique for root coverage. *J Periodontol.*, 56:715-20.
- Miller PD. Jr. 1985. A classification of marginal tissue recession. *Int J Periodontics Restorative Dent.*, 5:9-13.
- Pini Prato G., Pagliaro U., Baldi C., Nieri M., Saletta D., Cairo F. et al., 2000. Coronally advanced flap procedure for root coverage. Flap with tension versus flap without tension: A randomized controlled clinical study. *J Periodontol.*, 71:188-201.
- Pini Prato G., Tinti C., Vincenzi G., Magnani C., Cortellini P., Clauser C. 1992. Guided tissue regeneration versus mucogingival surgery in the treatment of human buccal gingival recession. *J Periodontol.*, 63:919-28.
- Rateitschak KH., Rateitschak EM., Wolf HF., Hassell TM. 1989. Color Atlas of Dental Medicine. Periodontology. Vol. 1. New York: Thieme Medical Publishers, Inc. p. 304.
- Romagna-Genon C. 2001. Comparative clinical study of guided tissue regeneration with a bioabsorbable bilayer collagen membrane and subepithelial connective tissue graft. *J Periodontol.*, 72:1258-64.
- Rosetti EP., Marcantonio RA., Rossa C. Jr, Chaves ES., Goissis G., Marcantonio E. Jr. 2000. Treatment of gingival recession: Comparative study between subepithelial connective tissue graft and guided tissue regeneration. *J Periodontol.*, 71:1441-7.
- Saletta D., Pini Prato G., Pagliaro U., Baldi C., Mauri M., Nieri M. 2001. Coronally advanced flap procedure: Is the interdental papilla a prognostic factor for root coverage? *J Periodontol.*, 72:760-6.
- Sullivan HC., Atkins JH. 1968. Free autogenous gingival grafts. I. Principles of successful grafting. *Periodontics.*, 6:121-9.
- Tal H., Moses O., Zohar R., Meir H., Nemcovsky C. 2002. Root coverage of advanced gingival recession: A comparative study between acellular dermal matrix allograft and subepithelial connective tissue grafts. *J Periodontol.*, 73:1405-11.
- Wei P., Laurell L., Geivelis M., Ligren MW., Maddalozzo D. 2000. Acellular dermal matrix allografts to achieve increased attached gingival. *J Periodontol.*, 71:297-305.
- Zucchelli G., De Sanctis M. 2000. Treatment of multiple recession-type defects in patients with esthetic demands. *J Periodontol.*, 71:1506-14.