



ISSN: 0975-833X

Available online at <http://www.journalcra.com>

INTERNATIONAL JOURNAL
OF CURRENT RESEARCH

International Journal of Current Research
Vol. 12, Issue, 08, pp.13186-13192, August, 2020

DOI: <https://doi.org/10.24941/ijcr.39371.08.2020>

RESEARCH ARTICLE

DIVERSITY OF FISHES IN BELES AND GILGEL BELES RIVERS, ABAY BASIN, ETHIOPIA

Zelege Berie^{1*}, Abebe Getahun² and Eshetie Dejen³

¹Gilgel Beles teacher's College, Department of Biology and Chemistry, Ethiopia

²Addis Ababa University, College of Natural Science, Ethiopia

³Amhara Regional Agricultural Research Institute, P.O. Box 794, Bahir Dar, Ethiopia

ARTICLE INFO

Article History:

Received 15th May, 2020
Received in revised form
21st June, 2020
Accepted 24th July, 2020
Published online 30th August, 2020

Key Words:

Rivers, Beles,
Gilgel Beles,
Diversity.

ABSTRACT

Diversity of fish species of Beles and Gilgel Beles Rivers were studied. Fishes were sampled using gill nets of various stretched mesh sizes, and hooks and lines. Identification of fish was made to species level by comparing the sample characters with taxonomic keys found in literatures and specimens deposited. Shannon diversity index (H') was used to evaluate species diversity of fishes. A total of 1124 individual fishes belonging to 5 orders, 7 families, 13 genera and 23 species were caught in Beles and Gilgel Beles Rivers. Cyprinidae was best-represented family with the highest number of fish species in the rivers while *Labeo* and *Labeobarbus* were the best-represented genera. A higher number of species were recorded in the Beles River (22 species) than in the Gilgel Beles River (5 species). The diversity index was higher in the Beles River ($H'=2.43$) than in the Gilgel Beles River ($H'=0.88$). In general, Beles River is richer than Gilgel Beles River, in terms of total numbers of fish caught and number of species recorded. In order to have a better knowledge of the fish populations in the study area, further studies are required on diversity, abundance and biology of the fishes. Therefore, sustainable utilization and conservation measures should be taken in the two rivers.

Copyright © 2020, Zelege Berie et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Zelege Berie, Abebe Getahun and Eshetie Dejen. 2020. "Diversity of fishes in beles and gilgel beles rivers, abay basin, Ethiopia", *International Journal of Current Research*, 12, (08), 13186-13192.

INTRODUCTION

Species diversity and abundance reflect the quantity and quality of the available habitat. The decline in abundance of freshwater fish in the world has been of concern for over one hundred years. Since the twentieth century, many fish species have suffered continuing declines in abundance and distribution, some at alarming levels. This includes many of the smaller species as well as all of the species targeted by inland commercial and recreational fisheries. These declines in abundance are commonly attributed to factors such as general habitat degradation (Llewellyn, 1983), modified patterns of stream flow (Gehrke et al., 1995), interrupted migratory pathways (Mallen-Cooper et al., 1995), reduced water quality and pollution (4), introduction of alien fish and diseases (Koehn, 1990), illegal fishing and commercial over fishing (Brown, 1992) and altered biotic interactions (Schiller, 1997). As in many parts of the world, population growth, agricultural development and industrialization contribute to the loss of species diversity of freshwater fishes in Ethiopia (Abebe Getahun and Stiassny, 1998).

Wide spread deforestation, degradation of the pristine environment, and other human induced factors might have left many Ethiopian streams, specially the northern ones, devoid of fish but the apparently resilient cyprinids (Abebe Getahun and Stiassny, 1998). As in many parts of the Ethiopia, human activities degrade fish habitat in numerous ways in study area. Wild fire, logging, impoundment, canalisation and agricultural activities are some of the major activities that degrade fish habitat. At present, we have no evidence of species extinction from Ethiopian freshwaters (Harrison and Stiassny, undated cited in (Abebe Getahun and Stiassny, 1998)) resulting from degradation of environment. One of the main reasons is a lack of definitive information on diversity of freshwater fishes and inconclusive data on the freshwater fish species. Beles and Gilgel Beles Rivers are flowing to the lower course of Abay in which adequate attention has not been given in the study of the diversity, abundance and economical potential of the fish fauna due to the presence of some inaccessible mountains and rugged geographical features. The absence of fishery data on these rivers triggers the researcher to conduct this study. Therefore, the study attempted to identify species composition of fishes in Beles and Gilgel Beles Rivers and evaluate the species diversity of Beles and Gilgel Beles Rivers fishes.

*Corresponding author: Zelege Berie,
Gilgel Beles teacher's College, Department of Biology and
Chemistry, Ethiopia.

MATERIALS AND METHODS

Site selection: A reconnaissance survey was conducted together with the research advisor to fix sampling sites. The survey was conducted in four sub areas along the Beles and Gilgel Beles Rivers. Two sampling sites were selected from each river taking into consideration the velocity of water, habitat type, altitude, depth of water, vicinity to road and substrate type.

Fieldwork: Three surveys were conducted to collect specimens from the sampling sites. The samples were taken in November, March and May. November and May were wet months while March was dry month. Gill nets with different mesh sizes were used to collect fishes. Multiple hooks and lines were used in areas where gill nets were not suitable. Immediately after retrieval, fishes were removed and total length and total weight of each specimen were measured. Total length was measured to the nearest 0.1 cm.

Laboratory studies: The specimens were soaked in tap water for a week to wash the formalin from the specimens. Then, they were transferred to 75 % ethanol. Identification was made to species level by comparing the sample characters with taxonomic keys found in the literature and specimens deposited at the Fisheries Laboratory, Department of Biology, AAU, and also at Bahir Dar fisheries and other aquatic life Research Center, and at National Fisheries and Other Living Aquatic Resources Research Centre, Sebeta. Keys found in (Shibru, 1973; Bishai, 1997; Boulenger, 1909; Boulenger, 1911; Boulenger, 1915; Boulenger, 1916; Lévêque, 1990; Lévêque, 1992; Eschmeyer, 2006; Nagelkerke, 1997) were used for identification. Meristic and morphometric characters were assessed for comparison purpose.

Data analysis: Generally, SPSS for Windows (version 10) and MINITAB (version 14) were used to perform the calculations and statistical analysis. Shannon diversity index (H') was used to evaluate species diversity of fishes. The Shannon index of diversity (H') is a measure of the number of species weighted by their relative abundances. The Shannon's diversity index explains both the variety and the relative abundance of species (Næsje et al., 2004). H' was calculated as:

$$H' = - \sum p_i \ln p_i$$

Where, p_i is the proportion of individuals found in the i th species. Shannon's diversity index (H') was used to indicate diversity at different sampling sites and/or rivers. A high value indicates high species diversity. Significance of differences in species diversity between sampling sites and/or rivers was tested using T-test.

RESULTS AND DISCUSSION

Fish species of Beles and Gilgel Beles Rivers: A total of 23 fish species were recorded during the present study from Beles and Gilgel Beles Rivers (Table 2). These fishes were represented by a single class Actinopterygii (ray-finned fishes), seven families and five orders (Table 2). The Cyprinidae, Bagridae and Characidae were the best-represented families with respect to numbers of species; with 11, 3 and 3 species, respectively (Table 2).

Labeo and Labeobarbus were the best-represented genera with numbers of species; with five and four species, respectively (Table 2). The freshwater fish fauna of Beles and Gilgel Beles Rivers contain a mixture of Nilo Sudanic (e.g. *B. docmak*, *B. bajad*, *H. forskalii*, *L. forskalii*, *M. kannume*, *S. serratus* and *S. schall*), highland East African (e.g. *L. intermedius*, *L. nedgia*, *C. gariepinus* and *O. niloticus*) and Endemic forms (e.g. *V. beso*). A higher number of species were recorded in the Beles River (22 species) than in the Gilgel Beles River (5 species) in the present study (Table 3). The number of fish species was highest at BAB and lowest at Mh sites (Table 3). Cyprinidae was the best-represented family with the highest number of fish species both in the Gilgel Beles and Beles Rivers. Although there was a pronounced disparity in species composition between Beles and Gilgel Beles Rivers, there was an overlap in *L. forskalii*, *L. nedgia*, *L. intermedius* and *O. niloticus* in the present study. However, *V. beso*, which is found in Gilgel Beles River, was not found in Beles River. Thus, large incidence of Gilgel Beles fishes in Beles River is due to drainage connection between the two rivers. Although several fish collections have been made from the Blue Nile system, few reports of collections are available from Gilgel Beles and Beles Rivers. Fish surveys which were carried out in the Gilgel Beles and Beles Rivers by (JERBE, 1999) and (JERBE, 2000) respectively were comparable to the present study. Compared to Sanja, WabiShebele and Angereb Rivers, Beles River harbors more diverse fish fauna. Sanja, WabiShebele and Angereb Rivers harbor 8, 13 and 19 fishes, respectively (Genanew Tesfaye, 2006; JERBE, 2004; JERBE, 2005). However, fish species diversity of Beles River comparable to Alvero River (31), a tributary of Baro, in the lowland Gambela region (JERBE, 2001).

The species list obtained in the present study reveal a difference in number of species between Beles and Gilgel Beles Rivers, as it was also shown by the species list of (JERBE, 1999; JERBE, 2000). However, there was some difference in species composition between the present study and that of (JERBE, 1999; JERBE, 2000). Most of the species recorded during the present study in the Gilgel Beles and Beles Rivers were also recorded by JERBE surveys. Although (JERBE, 2000) listed 25 fish species from Beles River, the following seven species were recorded during the present study but were not listed by (JERBE, 2000): H. forskalii, C. gariepinus, H. longifilis, B. nurse, A. occidentalis, R. loati and L. bynni. Some species that were found by (JERBE, 2000) from Beles River were not found by the present study. These were Mormyrops anguilloides Linnaeus 1758, Mormyrus caschive Linnaeus 1758, M. hasselquistii, Micralestes acutidens Daget 1957, Distichodus engycephalus, Chelaethiops bibie, Garra sp., Leptocypris niloticus, Schilbe mystus, Chiloglanis sp., and Tetraodon lineatus. (JERBE, 1999) listed four fish species from Gilgel Beles River. L. forskalii is recorded in the present study from Gilgel Beles River but not (JERBE, 1999). However, Garra sp. was recorded by (JERBE, 1999) from Gilgel Beles River but not in the present study.

Differences seen in the species composition between the present study and that of (JERBE, 1999; JERBE, 2000) might be due to differences in the sampling efficiency, habitats and seasons. The higher number of species recorded by JERBE might be attributed to the flexibility of their gears. In addition, many of their gears can be classified as active gears, in contrast to the gill nets and multiple hooks and lines of present study.

Table 1. Estimated distances from Gilgel Beles at Mender hullet, altitude and coordinates of sampling sites

| Site | Code | Distance from Mh | Elevation (a.s.l) | Coordinate (GPS) |
|-------------------------------|------|------------------|-------------------|------------------------------------|
| Gilgel Beles at Mender hullet | Mh | - | 1011 m | 11° 09' 53.5'' N; 36° 20' 39.3'' E |
| Gilgel Beles at College | Coll | 2 km | 1007 m | 11° 09' 35.1'' N; 36° 20' 008'' E |
| Beles at bridge | BB | 6 km | 994 m | 11° 11' 56.7'' N; 36° 19' 31.7'' E |
| Beles at Babizenda | BAB | 156 km | 596 m | 11° 07' 54.8'' N; 35° 28' 13.6'' E |




Fig.1. Map of Beles and Gilgel Beles Rivers (unscaled) [9].

Table 2. Fish species of Beles and Gilgel Beles Rivers

| Species name | Comm on name (Gunuz) | Family | Order |
|---------------------------------------|----------------------|------------|-------------------|
| <i>R. loati</i> Boulenger 1901 | Abella | Cyprinidae | Cypriniformes |
| <i>V. beso</i> | - | | |
| <i>L. niloticus</i> | Tsemebebella | | |
| <i>L. horie</i> | - | | |
| <i>L. coubie</i> | - | | |
| <i>L. forskalii</i> | Tseyä | | |
| <i>L. cylindricus</i> | - | | |
| <i>L. bynni</i> Boulenger 1911 | Goshe | | |
| <i>L. intermedius</i> | - | | |
| <i>L. nedgia</i> Rüppell 1836 | - | | |
| <i>L. degeni</i> Boulenger 1902 | - | | |
| <i>C. gariepinus</i> | - | Clariidae | Siluriformes |
| <i>H. longifilis</i> Valenciennes1840 | - | | |
| <i>B. bajad</i> | - | Bagridae | |
| <i>B. docmak</i> | - | | |
| <i>A. occidentalis</i> | Jajuna | | |
| <i>S. serratus</i> | - | Mochokidae | |
| <i>S. schall</i> | Buwa | | |
| <i>H. forskhalii</i> | - | Characidae | Characiformes |
| <i>B. macrolepidotus</i> | Yechacheya | | |
| <i>B. nurse</i> Rüppell 1832 | Lekwar | | |
| <i>M. kannume</i> | Bebela | Mormyridae | Osteoglossiformes |
| <i>O. niloticus</i> | Begebella | Cichlidae | Perciformes |

Table 3. Species of Beles and Gilgel Beles Rivers (+ = present; - absent).

| Species | Sampling sites | | | | Rivers | |
|--------------------------|----------------|------|----|-----|---------|-------|
| | Mh | Coll | BB | BAB | G.Beles | Beles |
| <i>R. loati</i> | - | - | + | + | - | + |
| <i>V. beso</i> | + | + | - | - | + | - |
| <i>L. niloticus</i> | - | - | - | + | - | + |
| <i>L. horie</i> | - | - | - | + | - | + |
| <i>L. coubie</i> | - | - | - | + | - | + |
| <i>L. forskalii</i> | + | + | + | + | + | + |
| <i>L. cylindricus</i> | - | - | + | + | - | + |
| <i>L. bynni</i> | - | - | + | + | - | + |
| <i>L. intermedius</i> | + | + | + | - | + | + |
| <i>L. nedgia</i> | + | + | + | + | + | + |
| <i>L. degeni</i> | - | - | + | + | - | + |
| <i>C. gariepinus</i> | - | - | - | + | - | + |
| <i>H. longifilis</i> | - | - | - | + | - | + |
| <i>B. bajad</i> | - | - | - | + | - | + |
| <i>B. docmak</i> | - | - | + | + | - | + |
| <i>A. occidentalis</i> | - | - | - | + | - | + |
| <i>S. serratus</i> | - | - | - | + | - | + |
| <i>S. schall</i> | - | - | + | + | - | + |
| <i>H. forskhali</i> | - | - | - | + | - | + |
| <i>B. macrolepidotus</i> | - | - | + | + | - | + |
| <i>B. nurse</i> | - | - | - | + | - | + |
| <i>M. kannume</i> | - | - | + | + | - | + |
| <i>O. niloticus</i> | - | + | - | + | + | + |

Table 4. Number of species (N) and diversity index (H') for total catch at sampling sites and rivers.

| | Sampling sites | | | | Rivers | |
|----|----------------|------|------|-----|--------------|-------|
| | Mh | Coll | BB | BAB | Gilgel Beles | Beles |
| H' | 1.08 | 0.77 | 1.82 | 2.3 | 0.88 | 2.42 |
| N | 4 | 5 | 12 | 21 | 5 | 22 |

Table 5. Mean number of species (N) and diversity indices (H') for fish caught in Beles and Gilgel Beles Rivers

| H'/N | River | Mean+ SD | t | df | Sig. |
|------|--------------|-----------|------|----|------|
| H' | Gilgel Beles | 0.73±0.2 | 4.71 | 9 | 0 |
| | Beles | 1.81±0.52 | | | |
| N | Gilgel Beles | 3.67±0.82 | 3.49 | 9 | 0.01 |
| | Beles | 11.2±5.26 | | | |

Table 6. Number of species (N) and diversity index (H') for fish caught during wet and dry seasons.

| River | Gilgel Beles | | Beles | | Beles and Gilgel Beles | |
|--------|--------------|------|-------|------|------------------------|------|
| | Dry | Wet | Dry | Wet | Dry | Wet |
| Season | | | | | | |
| H' | 0.81 | 0.66 | 2.48 | 2.21 | 2.29 | 1.99 |
| N | 4 | 5 | 19 | 17 | 22 | 18 |

Table 7. Mean number of fish species (N) and diversity index (H') during wet and dry seasons

| H'/N | Season | River | Mean + SD. | t | df | Sig. |
|------|--------|------------------|--------------|------|----|------|
| H' | Wet | Beles & G. Beles | 1.16 ± 0.59 | 0.39 | 9 | 0.7 |
| | Dry | Beles & G. Beles | 1.33 ± 0.88 | | | |
| N | Wet | Beles & G. Beles | 6.57 ± 4.04 | 0.42 | 9 | 0.68 |
| | Dry | Beles & G. Beles | 8 ± 7.44 | | | |
| H' | Wet | Gilgel Beles | 0.72 ± 0.24 | 5.82 | 5 | 0 |
| | Wet | Beles | 1.75 ± 0.21 | | | |
| N | Wet | Gilgel Beles | 3.75 ± 0.96 | 3.98 | 5 | 0.01 |
| | Wet | Beles | 10.33 ± 3.21 | | | |
| H' | Dry | Gilgel Beles | 0.76 ± 0.14 | 3.98 | 5 | 0.01 |
| | Dry | Beles | 1.91 ± 0.98 | | | |
| N | Dry | Gilgel Beles | 3.5 ± 0.71 | 5.82 | 5 | 0 |
| | Dry | Beles | 12.5 ± 9.19 | | | |

Compared to JERBE surveys, a wider range of habitats and months were sampled in the present study. This might be a reason for some species that caught in the Present study that were not reported by JERBE.

Species diversity, according to H' , was higher in the Beles River ($H' = 2.42$) than in the Gilgel Beles River ($H' = 0.88$) for the total catch (Table 4). Among sampling sites, species diversity was highest at Babizenda ($H' = 2.3$) and lowest at

College ($H' = 0.77$) (Table 4). The number of fish species ranged from 6 to 19 with a mean \pm SD of 11.2 ± 5.26 in Beles River and it ranged 3 to 4 with a mean \pm SD of 3.67 ± 0.82 in Gilgel Beles River. Shannon diversity index ranged 1.67 to 2.6 with a mean \pm SD of 1.81 ± 0.52 in Beles River and it ranged 0.46 to 1.04 with a mean \pm SD of 0.73 ± 0.2 in Gilgel Beles River. There was significant variation in both mean number of fish species and diversity index between Beles and Gilgel Beles Rivers ($P < 0.05$) (Table 5). Biodiversity patterns are directly and indirectly influenced by the geomorphology of riverine landscapes, which may be perceived as a nested hierarchy (Ward, 1998). The number of fish species in Beles and Gilgel Beles Rivers appear to be negatively correlated with altitude. The increase in number of fish species from Mh to BAB sampling sites coincide with decline in elevation. The main pattern documented in this study, is the occurrence of a distinct headwater fauna, and a sequential down stream shift in species composition. The decrease in number of fish species from lower to upper reaches were consistent with the studies carried out in other areas by (Nikolsky, 1963; Sydenham, 1977; Golubtsov, 2003). The increase in species number from up stream sites to down stream sites was associated with change in catchment area, canopy closure, substrate type, distance from source, depth and width of rivers (Toham Andre Kandem and Teugels, 1998). These variables reflect longitudinal gradient in the study area. Width of river was the most important variable that coincided with increase in species number from Mh sites to BAB sites. A total of 21 fish species found in Beles River at BAB sampling site with its mean river width of 56.5 ± 2.12 m while the lowest number of species (4) in Gilgel Beles River at Mh sampling site with its mean river width of 31 ± 2 m. This result is consistent with the studies carried out in other areas. In tropical area as (Angermeier, 1983) in Panama, (EDDS, 1993) in India and (Toham Andre Kandem and Teugels, 1998) in Cameroon found respectively a significant relationship between species number and width of the river, and species number and increasing gradient of depth. A total of 12 fish species found in Beles River at BB sampling site with its mean river depth of 6.3 ± 0.5 m while the lowest number of species (4) in Gilgel Beles River at Mh sampling site with its mean river width of 2.42 ± 0.28 m. A similar result using depth gradient (Sheldon, 1968), stream order or river position in the gradient (Beecher, 1988; Oberdorff 1992; Oberdorff, 1993) have also been reported for temperate rivers. In addition, canopy closure and diversity of substrate type (sand, gravel and large rocks) were also most probable environmental gradient explaining the spatial distribution of species in the sampling sites. Thus, the presence of year round dense vegetation and higher catchment area, diversity of substrate, river depth and width might contribute to high species diversity in Beles River than in Gilgel Beles.

Species diversity during wet and dry seasons: A higher number of species was recorded in the dry than in the wet season in total catch (22 versus 18 species) (Table 5). *R. loati*, *V. beso*, *L. niloticus*, *L. forskalii*, *L. horie*, *L. bynni*, *L. intermedius*, *L. nedgia*, *L. degeni*, *B. docmak*, *B. macrolepidotus*, *B. nurse*, *A. occidentalis*, *S. serratus*, *S. schall*, *M. kannume* and *O. niloticus* were found both during wet and dry seasons in total catch. *L. coubie*, *C. gariepinus*, *B. bajad*, *H. longifilis* and *H. forskalii* were collected during dry season where as *Labeo cylindricus* was collected during wet season. There was no pronounced disparity in the species composition between wet and dry seasons in Gilgel Beles River (Table 6). However, there was marked difference in species composition

during wet and dry seasons in Beles River (Table 6). The value of Shannon's diversity index was 0.81 and 0.66 in dry and wet seasons in Gilgel Beles River, respectively (Table 6). The index was also higher in the dry ($H' = 2.48$) than wet season ($H' = 2.21$) in Beles River (Table 6). Shannon's diversity index (H') indicated that species diversity was higher in dry than wet season in each river. The species diversity was also higher in the dry ($H' = 2.29$) than wet season ($H' = 1.99$) for the total catch (Table 6). The highest species diversity was obtained in Beles River during dry season ($H' = 2.48$) while the lowest in Gilgel Beles River during wet season ($H' = 0.66$) (Table 6). The number of fish species ranged from 3 to 14 with a mean \pm SD of 6.57 ± 4.04 in wet seasons in Beles and Gilgel Beles Rivers and it ranged 3 to 19 with a mean \pm SD of 8 ± 7.44 in dry season. Shannon diversity index ranged from 0.46 to 1.99 with a mean \pm SD of 1.16 ± 0.59 in wet season in Beles and Gilgel Beles Rivers and it ranged 0.75 to 2.6 with a mean \pm SD of 1.33 ± 0.88 in dry season. There was significant difference in the mean number of fish species and diversity index between Beles and Gilgel Beles Rivers both in wet and dry seasons ($P < 0.05$) (Table 7). However, there was no significant difference in the mean number of fish species and diversity indices between wet and dry seasons in the total catch ($P > 0.05$) (Table 7). There might be several reasons for changes in catches between wet and dry seasons. For example, variation in available habitats and gill net efficiency might contribute to variations in the catches. The higher number of species recorded during dry season than wet season attributed to a wider range of habitats sampled. This was mainly due to habitats suitable for gill net sampling during dry season. However, during wet season trees that grow hanging their branches down to the water on either side of the riverbank hinder reaching residency of fish in their habitats. Habitats targeted to gill nets were seldom deep water areas where some fish often reside dislocated by current of rivers during wet seasons. In addition, during wet season the efficiency of gill nets were decreased by logs, leaves, roots etc that were brought by flooding into rivers. Thus, differences seen in the data between wet and dry season may be due to differences in available habitats and gill net efficiency.

Conclusions and Recommendation

Conclusions

Beles River is richer than Gilgel Beles River, in terms of both total numbers of fish caught and numbers of species recorded. A higher number of species were recorded in the Beles River (22 species) than in the Gilgel Beles River (5 species) in the present study. The increase in number of fish species from lower to upper reaches of study area coincide with decline in elevation. The Cyprinidae, Bagridae and Characidae were the best-represented families with respect to numbers of species; with 11, 3 and 3 species, respectively. The Labeo and Labeobarbus were the best-represented genera with numbers of species. L. nedgia and L. degeni considered belonging to a single lip morphotype (Labeobarbus nedgia) endemic to Lake Tana by (41) are found in Beles River. The species diversity was also higher in the Beles River ($H' = 2.42$) than in the Gilgel Beles River ($H' = 0.88$) for total catch. A higher number of species was recorded in dry than wet season in total catch (22 versus 18 species). The species diversity was also higher in the dry ($H' = 2.29$) than wet season ($H' = 1.99$) for the total catch.

Recommendation

In order to have a better knowledge of the fish populations detailed studies and investigations are required on diversity and abundance of fish species in Abay basin in general and in Beles and Gilgel Beles Rivers in particular, especially at the lower reaches of Beles River. In addition, detailed knowledge on the biology and behaviour of most of the species are still lacking. Therefore, further studies are required on the biology and behaviour of fishes in the study area.

Acknowledgment

Lastly, we thank Graduate and Research Program of Addis Ababa University for financial support and Amhara Regional Agricultural Research Institute (ARARI) through its Bahir Dar FOALRC for their assistance all logistic necessities for fieldwork.

REFERENCES

- Abebe Getahun and Stiassny, M.L.J. 1998. The Fresh water Biodiversity crisis: the case of the Ethiopian fish fauna. *SINET: Ethiop. J. Sci.* 212 : 207-230.
- Angermeier, P.L. and Karr, J.R. 1983. Fish communities along environmental gradients in a system of tropical streams. *Environ. Biol. Fish* 9:117-135.
- Beecher, H.A., Dott, E.R. and Fernaw, R.F. 1988. Fish species richness and stream order in Washington streams. *Environ. Biol. Fish* 22: 193 - 209.
- Begon, M., Harper, J.L. and Townsend, C.R. 1990. Ecology: Individuals, populations and communities. 2nd ed. Black well Scientific Publications, 945 Pp.
- Bishai, M.H. and Khalil, T.M. 1997. Fresh water fishes of Egypt. Publication of National Biodiversity Unit. No. 9. 229 Pp.
- Boulenger, G.A. 1909. Catalogue of the fresh water fishes of Africa. Vol. I. *British Museum Natural History*, London, 373 Pp.
- Boulenger, G.A. 1911. Catalogue of the fresh water fishes of Africa. Vol. II. *British Museum Natural History*, London, 529 Pp.
- Boulenger, G.A. 1915. Catalogue of the fresh water fishes of Africa. Vol.III. *British Museum Natural History*, London, 526 Pp.
- Boulenger, G.A. 1916. Catalogue of the fresh water fishes of Africa. Vol.IV. *British Museum Natural History*, London, 392 Pp.
- Brown, P. 1992. The Murrumbidgee River fishery: a historical perspective. In: Roberts, J. and Oliver, R. eds.. *The Murrumbidgee, past and present*, 20-26. CSIRO, Griffith.
- EDDS, D.R. 1993. Fish assemblage structure and environmental correlates in Nepal's Gandaki River. *Copeia* 1: 48 - 60.
- Eschmeyer, W.N. 2006. *Catalog of fishes*. California Academy of Sciences Revised version of April 2006. <http://www.fishbase.org/search.php>
- FDROE, MOWR, Federal Democratic Republic of Ethiopia, Ministry of Water Resources 2000. Abay basin integrated development master plan project. Phase 2. Sectoral report. Vol.1.c.
- Gehrke, P.C., Brown, P., Schiller, C.B., Moffatt, D.B. and Bruce, A.M. 1995. River regulation and fish communities in the Murray - Darling River system, Australia. *Regulated Rivers: Research and Management* 11: 363 -375.
- Genanew Tesfaye 2006. Diversity, relative abundance and biology of fishes in Angereb and Sanja Rivers, Tekeze basin, Ethiopia. M. Sc.Thesis, Addis Ababa University, Ethiopia. 89 Pp.
- Golubstov, A.S., Darkov, A.A., Dgebuadze, Yu.Yu. and Mina, M.V. 1995. An Artificial key to fish species of the Gambela region The White Nile basin in the limits of Ethiopia. Joint Ethio-Russian Biological Expedition, Addis Ababa, Ethiopia. 84 Pp.
- Golubtsov, A. S., and Mina, M. V. 2003. Fish species diversity in the main drainage systems of Ethiopia: current state of knowledge and research perspectives. *Ethiop. J. Natu. Reso.* 2003. 52: 281-318.
- JERBE 1999. Report on the studies carried out by the freshwater biology group of the joint Ethio - Russian biological expedition.
- JERBE 2000. Report on the studies carried out by the freshwater biology group of the joint Ethio-Russian biological expedition.
- JERBE 2001. Report on the studies carried out by the freshwater biology group of the joint ethio-russian biological expedition.
- JERBE 2004. Report on the studies carried out by the freshwater biology group of the joint ethio-russian biological expedition.
- JERBE 2005. Report on the studies carried out by the freshwater biology group of the joint ethio-russian biological expedition.
- Koehn, J. and O'Connor, W. 1990. Threats to Victorian native freshwater fish. *Victorian Naturalist* 107: 5-12.
- Lévêque, C., Paugy, D. and Teugels, G, G. 1990. The Fresh and Brackish water fishes of West Africa. Paris, France Vol.1. 386 Pp.
- Lévêque, C., Paugy, D. and Teugels, G, G. 1992. The Fresh and Brackish water fishes of West Africa. Paris, France Vol. 2. 521 Pp.
- Llewellyn, L.C. 1983. The distribution of Fish in New South Wales. Australian Society for Limnology, Special Publication No.7, 23 pp.
- Mallen - Cooper, M. and Harris, J. H. 1993. Fish migration in the Murray Darling system and the decline of the Silver Perch. Australian Government Publishing Service, Canberra. 41 Pp.
- Mallen-Cooper, M., Stuart, I., Hides-Pearson, F. and Harris, J. 1995. Fish migration in the Murray River and assessment of the Torrumbarry fishway: New South Wales Fisheries Research Institute, Sydney.
- Næsje, T.F., Hay, C.J., Nickanor, N., Koekemoer, J.H., Strand, R., and Thorstad, E.B. 2004. Fish populations, gill net catches and gill net selectivity in the Kwando River, Namibia. Norwegian Institute for Nature Research Tungasletta 2, NO-7485 Trondheim, Norway. 65 Pp.
- Nagelkerke, L. A. J. 1997. The barbs of Lake Tana, Ethiopia – morphological diversity and its implication for taxonomy, trophic resource partitioning, and fisheries. Ph.D. thesis, Wageningen University.
- Nagelkerke, L. A. J. and Sibbing, F. A. 1996. A revision of the barbus *Barbus* spp., Cyprinidae, Teleostie of Lake Tana, Ethiopia, with a description of seven new species. Department of Experimental Animal Morphology and Cell Biology, Agriculture University, Marijkeweg40, 6709PG Wageningen, The Netherlands. 170 Pp.
- Nikolsky, G.V. 1963. *The Ecology of fishes*. Academic Press, London and New York. 352 Pp.

- Oberdorff, T and Porcher, J.P. 1992. Fish assemblage structure in Brittany streams France. *Aqua. Living Resour.*5:215 - 223.
- Oberdorff, T., Guibert, E. and Lucetta, J.U. 1993. Patterns of fish species richness in the seine river basin, France. *Hydrobiologia* 259: 157-169.
- Schiller, C.B., Bruce, A.M. and Gehrke, P.C. 1997. Distribution and abundance of native fish in New South Wales Rivers. In: Harris, J.H. and Gehrke, P.C. Eds. "*Fish and rivers in stress: The NSW rivers survey*", 71-102. New South Wales Fisheries Office of Conservation and the Cooperative Research Centre for Freshwater Ecology.
- Sheldon, A.L. 1968. Species diversity and longitudinal succession in stream fishes. *Ecology* 49:193-198.
- Shibru Tedla 1973. *Fresh water fishes of Ethiopia*. Haile Selassie University, Dept. of Biology, Addis Ababa, Ethiopia. 107 Pp.
- Sydenham, D.H.J. 1977. The qualitative composition and longitudinal zonation of the fish fauna of the River Ogun, Western Nigeria. *Revue de Zoologie Africanes* 91: 974-996.
- Toham Andre Kandem and Teugels, G. 1998. Diversity pattern of fish assemblages in the lower Ntem River basin Cameroon, with notes on potential effects of deforestation. *Arch. Hydrobiology* 1414: 421-446.
- Ward, J.V. 1998. Riverine landscapes: Biodiversity patterns, Disturbance Regimes, and Aquatic Conservation. *Biological Conservation* 83 3: 269 - 278.
