

Available online at http://www.journalcra.com

International Journal of Current Research Vol. 17, Issue, 05, pp.32701-32705, May, 2025 DOI: https://doi.org/10.24941/ijcr.48854.05.2025

### INTERNATIONAL JOURNAL OF CURRENT RESEARCH

# **REVIEW ARTICLE**

#### **EFFECT OF BIOSTIMULANTS ON WHEAT YIELD AND GRAIN QUALITY**

#### Ricardo Bemfica Steffen<sup>1\*</sup> and Gerusa Pauli Kist Steffen<sup>2</sup>

<sup>1</sup>Researcher, Dr., BioTec RS Tecnologia e Consultoria; <sup>2</sup>Researcher, Dr., Secretaria da Agricultura, Pecuária, Produção Sustentável e Irrigação, Rio Grande do Sul, Brazil (SEAPI)

ARTICLE INFO	ABSTRACT
<i>Article History:</i> Received 20 <sup>th</sup> February, 2025 Received in revised form 11 <sup>th</sup> March, 2025 Accepted 26 <sup>th</sup> April, 2025 Published online 28 <sup>th</sup> May, 2025	Wheat ( <i>Triticum aestivum</i> L.) is one of the most important agricultural crops worldwide, serving as both a staple food and a primary source of income for millions. Its grain quality is influenced by genetic, environmental, and management related factors, with test weight (hectoliter weight) being a key parameter for commercial classification. This study aimed to evaluate the effect of the biostimulant Penergetic, applied in different forms and dosages, on wheat yield and grain quality. The field experiment was conducted in Santa Maria. Bio Grande do Sul Brazil using the OPS Soberapo
Key words:	cultivar in a randomized block design with six treatments and eight replications. Treatments involving
Biological Fertility, Biostimulant, Sustainable Agriculture.	the combined application of the biostimulant in both liquid and solid formulations resulted in improved foliar health, indicating reduced biotic stress and enhanced biological balance in the soil. Consequently, significant increases in both grain yield and quality were observed, particularly in the
*Corresponding author	treatment with the highest dosage, which achieved a test weight exceeding 80 kg.hL <sup>-1</sup> . The biostimulant effect of the commercial Penergetic product contributed to enhanced enzyme activity in
Ricardo Bemfica Steffen	the soil, improved plant nutrient uptake, better foliar health, and overall vegetative development. The findings indicate that biostimulants based on Penergetic Technology significantly increase wheat yield and grain quality, with the most pronounced benefits observed under the combined application of both liquid and solid formulations at the highest tested dosage. These results emphasise the essential role of soil biological fertility in determining wheat crop performance.
Comministant Discourse Comming	essential role of soil biological fertility in determining wheat crop performance.

*Copyright*©2025, *Ricardo Bemfica Steffen and Gerusa Pauli Kist Steffen.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Citation: Ricardo Bemfica Steffen and Gerusa Pauli Kist Steffen, 2025.* "Effect of biostimulants on wheat yield and grain quality". International Journal of Current Research, 17, (05), 32701-32705.

# **INTRODUCTION**

Wheat cultivation represents one of the most important agricultural commodities of global relevance, playing a vital role in food security and the global economy. Wheat is a significant source of carbohydrates, proteins, and other essential nutrients, forming a dietary staple for billions of people across various regions of the world (Ulukan, 2024; Voučko et al., 2025). Wheat grains have widespread applications in the food industry, particularly in the production of bread, pasta, and other staple food products consumed by diverse populations across cultures and traditions (Bentley et al., 2022; Ulukam, 2024). Countries such as China, India, the United States, and Russia are prominent wheat producers, holding strategic positions in international trade and influencing both local economies and the global dynamics of supply and demand. The growing demand for wheat, driven by population growth and changing dietary patterns, makes the sustainability of wheat production a central issue for ensuring economic and food stability on a global scale (Abys et al., 2024). The quality of wheat grains is a determining factor in the production of high-value food products and is influenced by a range of genetic, agronomic, and environmental factors. It is characterized by parameters such as protein content, gluten levels, color, grain integrity, and test weight (hectoliter

weight), all of which directly affect its suitability for the food industry (Ma et al., 2021). Among these, test weight is the most used parameter in grain commercialization between farmers and the food industry (Kumar et al., 2024). Factors such as climate, fertilization, and field management practices significantly impact grain quality, potentially leading to variations that affect both yield and the technological properties of wheat. Microbial activity in the soil and the resulting biological fertility of the soil-plant system play a decisive role in determining wheat test weight. Test weight is directly related to the density and integrity of the grains, reflecting their physical and sanitary characteristics. Soils with high biological fertility, defined by rich microbial biodiversity, elevated organic matter content, and a well-structured soil profile, promote optimal root development in plants, leading to grains with greater mass and density (Hebe et al., 2022; Lamlom et al., 2023). In this context, the presence of beneficial microorganisms, such as bacteria and fungi, enhances the availability of essential nutrients for wheat, improves soil water retention, and increases the plants' environmental resilience to stressors. Consequently, agricultural practices that foster soil biological health represent effective strategies for enhancing both wheat yield and grain quality. Among the products based on technological

innovations with the effect of reestablishing balance in agricultural systems and increase of the crops productivity, the Penergetic technology has attracted the attention of farmers and the scientific community. Research works have already demonstrated the effect of the product on agricultural crops and on the biological activity of soils in different soils around the world (Kadziuliene *et al.*, 2005; Jakiene *et al.*, 2008; Jankauskiene & Surviliene, 2009; Cobucci *et al.*, 2015; Souza *et al.*, 2017; Franco Júnior *et al.*, 2018; Franco Júnior *et al.*, 2019). This study aimed to evaluate the effectiveness of the biostimulant Penergetic in improving wheat productivity and grain quality under different application forms and dosages.

## MATERIALS AND METHODS

The experiment was conducted under field conditions at the experimental area of Biomonte Research and Development, located in the municipality of Santa Maria, RS, Brazil, with the following field coordinates: 29°38'48.1"S and 53°57'40.7"W. The experimental area has soil characterized as Dystrophic Red Argisol. The field experiment was carried out from June 25, 2024, to October 23, 2024. The ORS Soberano wheat cultivar, known for its medium cycle and high quality, was used. The crop was established using a no-tillage system, with sowing on June 25, 2024, and emergence on July 2, 2024. The experimental design used was a randomized block design, with 6 treatments and 8 replications. The experimental plots measured 20 meters in width by 50 meters in length, totaling 1000 m<sup>2</sup>. For evaluation purposes, a central useful area of 300 m<sup>2</sup> was considered. The sowing was performed with a 17 cm row spacing and a seed density of 140 kg per hectare.

Different combinations of the biostimulant Penergetic were applied in liquid form (molasses) and solid form (sikron). The evaluated treatments were: T1: Control treatment (no Penergetic applied); T2: 500g of Penergetic b (sikron) and 500g of Penergetic p (sikron); T3: 1000mL of Penergetic b (molasses) and 500mL of Penergetic p (molasses); T4: 1000mL of Penergetic b (molasses) and 500g of Penergetic p (sikron); T5: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses); T6: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses) + 300g of Penergetic p (sikron). The Penergetic products in solid form were previously mixed with water and added to the spray solution. The Penergetic b products (liquid and solid) were applied 15 days before wheat sowing. The Penergetic p products were applied at the BBCH 13-14 growth stage (Figure 1).



Figure 1. Application scheme of the biostimulant Penergetic in wheat crops

In all treatments, standard fertilization and phytosanitary management practices for wheat in the southern region of Brazil were used. Treatments T2 through T6 were conducted in plots that had been using the Penergetic technology for 2 years.

Potassium and phosphate fertilization were omitted, with only nitrogen fertilization (urea) applied at a rate of 200 kg/ha. The applications were carried out on July 13, 2024, and July 16, 2024. Phytosanitary management followed the technical recommendations for wheat cultivation in the southern region of Brazil. Wheat yield was evaluated by grain weight per area, with moisture content adjusted to 12%, and grain quality was determined according to the methodology used by Awulachew (2020). Data were submitted to analysis of variance, with means compared by Tukey's test at a 5% probability of error using statistical software.

## **RESULTS AND DISCUSSION**

The use of the biostimulant Penergetic resulted in an increase in microbial activity in the soil, as shown in the bioanalysis presented in Figure 2. Microbial activity in the soil was significantly higher in treatments T5 and T6, where higher dosages of the biostimulant were used in the combination of liquid and solid forms. According to Akbar et al. (2025) and Semenov et al. (2025), microbial activity in the soil is a key factor for high-potential regenerative agriculture. The authors suggest that management practices and technologies that increase microbial activity in the soil are essential tools for modern agriculture. The increases in microbial activity in the soil likely resulted in significant differences in plant development in the field, particularly in terms of the health of the wheat's above-ground parts. In treatments where higher dosages of the biostimulant Penergetic were applied (combination of liquid and solid products), the wheat leaves did not show any visible signs of biotic stress during the ear formation stage (Figure 3).



Figure 2.  $\beta$ -glucosidase enzyme activity in the soil under different treatments. Treatments:T1: control;T2: 500g of Penergetic b (sikron) and 500g of Penergetic p (sikron);T3: 1000mL of Penergetic b (molasses) and 500mL of Penergetic p (molasses); T4: 1000mL of Penergetic b (molasses) and 500g of Penergetic p (sikron); T5: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses);T6: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses) + 300g of Penergetic p (sikron).

As a microbial stimulator, Penergetic directly enhances microbial activity in the soil, promoting biological balance (Steffen *et al.*, 2020; Hata *et al.*, 2022). Biological balance in the soil is crucial for plant health as it fosters the interaction between beneficial organisms and the plant root system, resulting in more structured soil with higher biological fertility, better water and nutrient retention capacity, and enhanced plant development and resistance to diseases. It is likely that the improvement in plant health contributed to the increased productivity and quality of wheat grains (Figures 3 and 4). The

better the biological fertility of the soil, the better the conditions for maintaining microbial communities that interact with the plant root system, promoting growth and/or plant protection.



Figure 3. Wheat leaves during the ear formation stage in the different evaluated treatments.Treatments:T1: control;T2: 500g of Penergetic b (sikron) and 500g of Penergetic p (sikron);T3: 1000mL of Penergetic b (molasses) and 500mL of Penergetic p (molasses);T4: 1000mL of Penergetic b (molasses) and 500g of Penergetic p (sikron);T5: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses);T6: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses);T6: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses) + 300g of Penergetic p (sikron).

The establishment of symbioses between plants and microorganisms, as well as between different groups of microorganisms within microbial communities, results in an increase in the diversity of microbial metabolites (acids and organic compounds, hormones, amino acids, peptides, and polysaccharides) that regulate the physiological and biochemical processes of microorganisms and plants in symbiosis (Singh et al., 2024). The direct outcome of these interactions is the improved ability of plants to reach their maximum productive potential and produce higher-quality grains. In this study, all treatments that received the application of the biostimulant Penergetic showed higher wheat productivity compared to the control treatment. The treatments with higher dosages of the biostimulant (through the combination of liquid and solid product applications) resulted in the highest productivity averages. Treatment T6 (1000mL of Penergetic b molasses + 500g of Penergetic b sikron and 500mL of Penergetic p molasses + 300g of Penergetic p sikron) increased wheat grain productivity by 74.9% compared to the control treatment (Figure 4). The average wheat yield in Rio Grande do Sul (Southern Brazil) was 2,839 kg per hectare in the 2024 growing season. Based on this information, it is evident that the average yield achieved with the use of the Penergetic biostimulant holds technical relevance for enhancing grain productivity. The treatments that resulted in higher wheat productivity also exhibited higher grain quality, a parameter evaluated by the size, shape, and uniformity of the grains (Figure 5). The quality of wheat grains is a crucial factor in determining their suitability for different uses, such as flour for baking, food products, or animal feed. Grain quality is influenced by various factors, including climatic conditions, agricultural practices, the genetic characteristics of each cultivar, and, directly, biological quality. The benefits of using the biostimulant Penergetic for wheat productivity and quality have already been studied by researchers from different research institutions (Artyszak & Gozdowski, 2021: Klevinskaitė, 2022; Pekarskas et al., 2011; Pekarskas, 2012; Pekarskas & Sinkevičienė, 2015; Pekarskas et al., 2017; Veverskytė, 2020; Steponavičienė et al., 2021).



Figure 4. Wheat productivity under the different evaluated treatments.Treatments:T1: control;T2: 500g of Penergetic b (sikron) and 500g of Penergetic p (sikron);T3: 1000mL of Penergetic b (molasses) and 500mL of Penergetic p (molasses);T4: 1000mL of Penergetic b (molasses) and 500g of Penergetic p (sikron);T5: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses);T6: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic b (sikron).

The effect of using the biostimulant Penergetic was observed for the hectoliter weight variable (Figure 6). The grains from plants that received the highest doses from the combination of sikron and molasses products showed higher quality of the final product, with hectoliter weight exceeding 80 (Figure 6). This result is very important as it is directly related to the price paid to the producer for the grain to be used in the food industry. Grains with higher hectoliter weight result in flour with better technical properties, such as greater water absorption capacity and higher elasticity, which are essential for bread and other product quality.



Figure 5. Wheat grains obtained from the different evaluated treatments.Treatments:T1: control;T2: 500g of Penergetic b (sikron) and 500g of Penergetic p (sikron);T3: 1000mL of Penergetic b (molasses) and 500mL of Penergetic p (molasses);T4: 1000mL of Penergetic b (molasses) and 500g of Penergetic p (sikron);T5: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses);T6: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses) + 300g of Penergetic p (sikron)

Additionally, higher-quality wheat exhibits greater uniformity and lower risk of contamination by mycotoxins, which adds value to the production, ensuring greater efficiency in the industrial process and better final quality for the consumer. These factors justify the higher payment by the industry, as higher-quality wheat contributes to a superior final product with better yield and less waste during production. The correlation between the biological fertility of the soil and the productive efficiency of the wheat crop was clearly demonstrated in this study. The data obtained reinforce the



Figure 6. Hectoliter weight obtained from wheat grains under the different evaluated treatments. Treatments:T1: control;T2: 500g of Penergetic b (sikron) and 500g of Penergetic p (sikron);T3: 1000mL of Penergetic b (molasses) and 500mL of Penergetic p (molasses);T4: 1000mL of Penergetic b (molasses) and 500g of Penergetic p (sikron);T5: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses);T6: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses);T6: 1000mL of Penergetic b (molasses) + 500g of Penergetic b (sikron) and 500mL of Penergetic p (molasses) + 300g of Penergetic p (sikron).

importance of agricultural practices and technologies that promote the enhancement of soil biological activity as an effective strategy to increase the quality and productivity of agricultural crops.

## CONCLUSION

Increases in productivity and quality indices for the wheat crop were achieved with the use of biostimulants from Penergetic technology. The dosage of the biostimulants and the nature of the commercial products influenced the biological activity of the soil, as well as the productivity and quality of the wheat grains. The combined application of liquid and solid products, along with higher dosages, resulted in greater microbial and productive bio-stimulation. Plants that received the biostimulants exhibited greater health, with direct positive impacts on yield and the qualitative characteristics of the grains. Among the parameters evaluated, hectoliter weight stood out as an indicator of superior quality, reflecting higher commercial value and better industrial performance of the grains. The treatment with the combined application of the biostimulant in both liquid and solid forms at the highest evaluated dosage (1000mL of Penergetic b molasses + 500g of Penergetic b sikron and 500mL of Penergetic p molasses + 300g of Penergetic p sikron) was the most efficient, providing the best results in productivity and quality.

#### STATEMENTS AND DECLARATIONS

**Funding:** The authors declare that no funds, grants, or other support were receivedduring the preparation of this manuscript.

**Competing Interests:** The authors declare that there is no conflict of interest of any kind involved incarrying out his scientific research study. Thus, the authors have no competing interests to declare that are relevant to the content of this manuscript.

**Data Availability:** The datasets generated during the current study are available from the corresponding author on reasonable request.

## REFERENCES

- Abys, C.;Skakun, S.; Becker-Reshef, I. (2024). Two decades of winter wheat expansion and intensification in Russia. *Remote Sensing Applications: Society and Environment*, 33, 101097.https://doi.org/10.1016/j.rsase.2023.101097
- Akbar, A. et al. (2025). Microbial diversity and its role in enhancing soil fertility: a comprehensive review. *The Research of Medical Science Review*, 3(2), p. 238-253.
- Artyszak, A.;Gozdowski, D. (2021). Application of growth activators and Plant Growth-Promoting Rhizobacteria as a method of introducing a "farm to fork" strategy in crop management of winter oilseed. *Sustainability*, *13*(6), p. 3562, 2021.https://doi.org/10.3390/su13063562
- Awulachew, M. T. (2020). Understanding basics of wheat grain and flour quality. *Journal of Health and Environmental Research*, 6(1), p. 10-26.https://doi.org/10.11648/j.jher.20200601.12
- Bentley, A. R. et al. (2022). Near-to long-term measures to stabilize global wheat supplies and food security. *Nature Food*, *3*(7), p.483-486. https://doi.org/10.1038/s43016-022-00559-y
- Cobucci, T.; Nascente, A. S.; Lima, D. P. (2015). Phosphate fertilization and Penergetic application in the yield of common bean. *Revista Agrariam*, 8(3), p.358-368. https://doi.org/10.33448/rsd-v11i11.34159
- Franco Junior, K. S. et al. (2018). Effect of cover crops and bioactivators in coffee production. *Coffee Science*, 13(4), p. 559 – 567.
- Franco Junior, K. S. et al. (2019). Influence of the use coverage plants and the bioactivator in the physical-biological characteristics of soil cultivated with coffee. *Coffee Science*, *14*(1), p. 116-122.
- Hata, F. T. et al. (2021). Bokashi, boiled manure and Penergetic applications increased agronomic production variables and may enhance powdery mildew severity of organic tomato plants. *Horticulturae*, 7(2), p. 27. https://doi.org/10.3390/horticulturae7020027
- Hebe, K. et al. (2022). Improve wheat productivity by using a combination of mineral nitrogen, organic and biological fertilizers under new sandy soil conditions. *Pakistan Journal of Biotechnology*, *19*(1), p. 13-22.https://doi.org/10.34016/pjbt.2022.19.1.13
- Jakiene, E.; Venskutonis, V.; Michevicius, V. (2008). The effect of additional fertilization with liquid complex fertilizers and growth regulators on potato productivity. *Sodininkysteir Darzininkyste*, 27(2), p.259-267.
- Jankauskiene, J.; Surviliene, E. (2009). Influence of growth regulator on seed germination energy and biometrical parameter of vegetables. *SodininkysteirDarzininkyste*, 28(3), p.69-77.
- Kadziuliene, Z. et al. (2005). Peculiarities of some legumes and cereals under organic farming system. *NJF Report*, *I*(1), p.103-106.
- Klevinskaitė, R. (2022). Biologiniųpreparatųiržem ėsdirbimote chnologijosįtakadirvožemiosavybėmsvasariniųkviečiųpas ėlyje. *In*: Conference Young Scientist 2023. 2022. p. 601-605.
- Kumar, P. et al. (2024). Impact of fertilizer levels and treated sewage water on heavy metal uptake, growth, agrometeorological indices, and quality of wheat. *Indian Journal of Agronomy*, *69*(3), p. 257-262. https://doi.org/10.59797/ija.v69i3.5519

- Lamlom, S. F.; Irshad, A.; Mosa, W. F. (2023). The biological and biochemical composition of wheat (*Triticum* aestivum) as affected by the bio and organic fertilizers. BMC plant biology, 23(1), 111.https:// doi.org/ 10.1186/s12870-023-04120-2
- Ma, M. et al. (2021). Current situation and key parameters for improving wheat quality in China. Frontiers in Plant Science, 12, 638525.https://doi.org/10.3389/fpls.2021.638525
- Pekarskas, J. (2012). Influence of growth activator Penergeticp on yield and chemical composition of barley in organic farm. *Žemėsūkio Mokslai*, 19(4), p. 249-256. https://doi.org/http://dx.doi.org/10.6001/zemesukiomoksl ai.v19i4.2586
- Pekarskas, J. et al. (2011). Effect of organic nitrogen fertilizers Provita and fermentator Penergetic-k winter wheat and on soilquality.*In*:

Environmental engineering: 8th international conference, May 19-20, 2011. *Vilnius, Lithuania: selected papers. Vol. 1: Environmental protection. Vilnius: Technika.* 

- Pekarskas, J.; Gavenauskas, A.; Dautartė, A. (2017). The effect of biological preparation Fertenat on spring wheat productivity and quality. Žemėsūkiomokslai, 24(1), p. 10-19.https://doi.org/10.6001/zemesukiomokslai.v24i1.3441
- Pekarskas, J.;Sinkevičienė, J. (2015). Effect of biopreparations on seed germination and fungal contamination of winter wheat. *Biologija*, 61(1), p. 25-33. https://doi.org/10.6001/biologija.v61i1.3108
- Semenov, M. V. et al. (2025). Microbiological indicators for assessing the effects of agricultural practices on soil health: A Review. Agronomy, 15(2), 335. https://doi.org/10.3390/agronomy15020335

- Singh, N. K. et al. (2024). Building soil health and fertility through organic amendments and practices: a review. *Asian Journal of Soil Science and Plant Nutrition*, 10(1), p. 175-197. https://doi.org/10.9734/ajsspn/2024/v10i1224
- Souza, A. A. de.; Almeida, F. Z. de.; Alberton, O. (2017). Growth and yield of soybean with Penergetic application. *Revista Scientia Agraria*, 18(4), p. 95-98.
- Steffen, R. B.; Steffen, G. P. K.; Maldaner, J. (2020). In Vitro activation of microbial growth in low frequency electromagnetic fields. *Journal of Agriculture and Environmental Sciences*, 9(1), p. 1-7.https://doi.org/10.15640/jaes.v9n1a1
- Steponavičienė, V. et al. (2021). The influence of biological preparations and their mixtures on soil agrochemical properties in winter wheat. In: Proceedings of the International Scientific Conference Rural Development. pp. 38-42.https://doi.org/10.15544/RD.2021.008
- Ulukan, H. (2024). Wheat production trends and research priorities: A global perspective. In Advances in Wheat Breeding: Towards Climate Resilience and Nutrient Security (pp. 1-22). Singapore: Springer Nature Singapore.
- Veverskytė, A. (2020). Biologiniųpre paratųirjųder iniųįtaka žiemi niųkviečiųligotumui. Organizaciniskomitetas, p. 96-99. https://doi.org/10.7220/1822-9913.2020
- Voučko, B. et al. (2025). Wheat: from nutrition to cultivation and technology. *Frontiers in Nutrition*, 12, 1563397.https://doi.org/10.3389/fnut.2025.1563397

\*\*\*\*\*\*